VIEQUES

Dr. Arturo Massol Deyá



Departamento de Biología Universidad de Puerto Rico Mayagüez, PR 00681-9012





Elements of a Bomb

Acetone

- * Acetyl Triethyl Citrate
- * Aluminum Powder
- * Ammonium Nitrate

Antimony

Barium

- * Boron
- * Butyl Stearate

Cadmium

Chromium

* Diazodinitrophenol

Diethyl Phthalate

Dimethyl Phthalate

Di-n-butyl phthalate

Di-n-octyl Phthalate

* Dichromated Aluminum Powder

- 2, 4- Dinitrotoluene
- 2, 6-Dinitrotoluene

Diphenylamine

- * Ethyl Centralite
- * HMX

Lead

* Magnesium Powder

Magnesium Alloy Powder

Mercury

Nickel Powder

Nitrocellulose

- * Nitroglycerin
- * Nitroguanidine
- * Nitrostarch
- * Pentolite
- * Pentaerythritol Tetranitrate (PETN)
- * Phosphorous

- * Potassium Nitrate
- * RDX (Cyclonite)
- * Resorcinol Selenium
- * Sodium Nitrate
- * Strontium Nitrate
- * Tetracene
- * Tetryl
- * Titanium Powder
- * TNR (Trinitroresorcinol)
- * TNT(Trinitotoluene)
- * Tri Amino Guanidine Nitrate

Vinyl Acetate

Vinyl Chloride


Zinc

* Zirconium Powder












Dispersion of natural trace elements is increased due to military activities.

Is pollution at the Atlantic Fleet Weapons Training Facility contained?









Why PLANTS?

- ✓ Transitory receptors of trace elements.
- ✓ Plants can concentrate environmental toxins, thus they can be good bioindicators of environmental degradation.
- ✓ Ecological significance: start-up of the food chain.
- ✓ Plants are relatively "immune" to human toxins.


 (no diabetes, hypertension or cancer cases have been reported in plants)


Potential use in clean up efforts: Phytoremediation



Methodology

- ✓ Life vegetation samples were manually collected.
- ✓ Each station was randomly sampled in a 2 to 5 m² circular plot. Sampling sites were independently established for each species. Vegetation analysis always included leaves material. Other plant components were selected as indicated. The sampling was consistently performed by selecting only large, green, healthy looking leaves. Samples from each species consisted of over 30 leaves picked alternately from upper, middle, and the lower sections of 5 to 10 individual plants.
- ✓ Samples were acid digested (2-4 replicates) and analyzed in an atomic absorption spectrophotometer.







Zonas Bajo Estudio por Massol y Díaz. CasaPueblo, Adjuntas PR.


Areas de Muestreo


- (a) Polígono de Prácticas Militares, Vieques
- **b** Zona Civil, Vieques
- (c) Antigua Zona de Prácticas Militares, Culebra
- d Bosque Xerofítico de Guánica
- (e) Fincas, Mayagüez y Las Marías
- (f) Zona Industrial, Peñuelas











...analysis revealed sea around Vieques is free of heavy metal contamination!

However, at that time, live ammunition wasn't being used.

ammunition wasn't being used.

Government Documentation

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY REGION II

IN THE MATTER OF:

United States
Department of the Navy

Atlantic Fleet Weapons Training Facility Viegues Island, PR

EPA ID Number PRD980536221

Respondent

ADMINISTRATIVE ORDER
ON CONSENT

DOCKET No. RCRA-02-2000-7301

Proceeding under Section 3008(h), of the Resource Conservation and Recovery Act, as amended.

In 1994, the EPA identified 102
 violations to water quality parameters.

The EPA Discharge Monitoring Reports identified excessive concentrations of the following contaminants in the coastal waters of Vieques:

Boron

Grease

Phenois

Selenium

Silver

Sulfates

Zinc

Cyanide (10x)

Cadmium (240x)

Chromium (13x)

Iron

Lead (105x)

Manganese


Mercury (4.6x)

Arsenic (6.6x)

Nitrogen (188x)

	Chemical Element (µg/g DW)						
	Pb	Co	Ni	Al	As	Cd	Cu
AFWTF-2001(Active)	33.32 (10.77)	29.60 (5.51)	2866 (1.58)	-	-	0.28 (0.13)	30.48 (3.63)
AFWTF-2004(Non-active)	8.14 (3.15)	10.61 (4.21)	3.43 (2.13)	1543 (67.7)	0.61 (0.46)	0.15 (0.15)	17.42 (1.83)
Guţnica-2001	5.58 (1.90)	4.19 (0.06)	1464 (4.75)	-	-	0.28 (0.01)	15.39 (4.16)
Guţnica-20032004	2.33 (2.24)	1.82 (0.43)	3.75 (1.85)	3412 (112.9)	1.04 (0.25)	0.28 (0.22)	1216 (2.83)





Pigeon Peas

Leaf samples (June 2000)

	Cu	Mn	Zn	Pb	Ni	Cd	Со
Monte Carmelo/Vieques	18.30	182.99	34.77	10.98	10.37	2.44	14.64
	12.58	203.73	27.56	10.19	10.19	3.00	14.98
average	15.44	193.36	31.17	10.59	10.28	2.72	14.81
std dev	(4.04)	(14.67)	(5.10)	(0.56)	(0.13)	(0.40)	(0.24)
Sector Gobeo/Vieques	28.87	77.63	87.57	54.43	7.10	0.95	7.10
	22.75	71.56	71.56	43.85	6.20	1.24	4.96
average	25.81	74.60	79.57	49.14	6.65	1.10	6.03
std dev	(4.33)	(4.29)	(11.32)	(7.48)	(0.64)	(0.21)	(1.51)
CONTROL	10.45	142.98	31.62	4.95	7.97	0.82	4.40
Cerro Las	11.33	132.69	29.58	4.98	7.46	0.83	3.87
Mesas/Mayagüez	11.08	130.69	30.4	4.55	7.67	0.85	4.55
	11.30	143.36	31.98	5.24	6.66	1.10	4.41
promedio	11.04	137.43	30.90	4.93	7.44	0.90	4.31
std dev	(0.41)	(6.68)	(1.11)	(0.28)	(0.56)	(0.13)	(0.30)
MC VIEQ/CONTROL	1.4	1.4	1.0	2.1	1.4	3.0	3.4
SG VIEQ/CONTROL	2.3	0.5	2.6	10.0	0.9	1.2	1.4



Monte Carmelo, Vieques (Puerto Rico) Pigeon Peas

Cu	Zn	Ni	Pb	Cd	Co
15.00	39.15	2.45	14.70	2.58	16.78
6.43	25.08	2.06	15.53	2.26	18.13
10.72	32.12	2.26	15.12	2.42	17.46
(6.06)	(9.95)	(0.28)	(0.59)	(0.23)	(0.95)
7.02	35.31	5.33	7.80	2.21	23.29
9.41	32.68	4.27	4.89	4.13	28.21
8.22	34.00	4.80	6.35	3.17	25.75
(1.69)	(1.86)	(0.75)	(2.06)	(1.36)	(3.48)
8.0	1.1	2.1	0.4	1.3	1.5
	15.00 6.43 10.72 (6.06) 7.02 9.41 8.22 (1.69)	15.00 39.15 6.43 25.08 10.72 32.12 (6.06) (9.95) 7.02 35.31 9.41 32.68 8.22 34.00 (1.69) (1.86)	15.00 39.15 2.45 6.43 25.08 2.06 10.72 32.12 2.26 (6.06) (9.95) (0.28) 7.02 35.31 5.33 9.41 32.68 4.27 8.22 34.00 4.80 (1.69) (1.86) (0.75)	15.00 39.15 2.45 14.70 6.43 25.08 2.06 15.53 10.72 32.12 2.26 15.12 (6.06) (9.95) (0.28) (0.59) 7.02 35.31 5.33 7.80 9.41 32.68 4.27 4.89 8.22 34.00 4.80 6.35 (1.69) (1.86) (0.75) (2.06)	15.00 39.15 2.45 14.70 2.58 6.43 25.08 2.06 15.53 2.26 10.72 32.12 2.26 15.12 2.42 (6.06) (9.95) (0.28) (0.59) (0.23) 7.02 35.31 5.33 7.80 2.21 9.41 32.68 4.27 4.89 4.13 8.22 34.00 4.80 6.35 3.17 (1.69) (1.86) (0.75) (2.06) (1.36)


August 2001


2ND Draft

Preliminary Evaluation of Health Risks Related to Naval Activities at the Atlantic Fleet Weapons Training Facility on Vieques, Puerto Rico

August 2000

"Bombing activities that occur on the eastern portion of the island are likely to produce airborne emissions that would be carried in a westerly direction." US Navy









QuickTime™ and a TIFF (LZW) decompressor are needed to see this picture.

Vieques / 2003-05 Departamento de Salud de PR 18°20'0" 18*20'0" 18°20'0" 18°0' -65°40'0

Trace Element Composition in Forage Samples from a Military Target Range, Three Agricultural Areas, and One Natural Area in Puerto Rico

ELBA DÍAZ AND ARTURO MASSOL-DEYA

Department of Biology, University of Puerto Rico, Mayagüez, Puerto Rico 00681 amassol@stahl.uprm.edu

ABSTRACT scarce. Trace Facilities (Al State Forest. with acid. Consideration of the Rico. Simila Sporobolus and marine

QuickTime[™] and a TIFF (LZW) decompressor are needed to see this picture.





Agradecimientos



- »Elba Díaz
- »Dustin Pérez
- »Ernie Pérez
- »Manuel Berríos
- Botánicos [UPR-Mayagüez]
- Familia Zenón-Encarnación
- Programa de Biotecnología Industrial, UPRM
- Casa Pueblo, Adjuntas
- Departamento de Salud de PR
- Escuela Elemental Playa Grande, Victor Belardo y los estudiantes de programa ambiental