



NAVAL FACILITIES ENGINEERING COMMAND
Washington, DC 20374-5065

NFESC User's Guide UG-2041-ENV

Guide for Incorporating Bioavailability Adjustments into Human Health and Ecological Risk Assessments at U. S. Navy and Marine Corps Facilities

Part 1: Overview of Metals Bioavailability

Prepared for

Naval Facilities Engineering Service Center
Port Hueneme, CA 93043-4370

and

Naval Facilities Engineering Command
Engineering Field Activity West
San Bruno, CA 94066-5006

Prepared by

Battelle
and
Exponent

July 2000

Approved for public release; distribution is unlimited.

ACKNOWLEDGEMENTS

The Navy and Battelle would like to acknowledge the following Navy Divisions, Offices, and Activities for providing review and suggestions for improving this document:

Naval Facilities Engineering Service Center, Port Hueneme, CA
Engineering Field Activity West, San Bruno, CA
Northern Division Naval Facilities Engineering Command, Lester, PA
Southern Division Naval Facilities Engineering Command, North Charleston, SC
Southwest Division Naval Facilities Engineering Command, San Diego, CA
Engineering Field Activity Northwest, Poulsbo, WA
Naval Environmental Health Center, Norfolk, VA
Office of the Chief of Naval Operations, Arlington, VA
Naval Surface Warfare Center, Dahlgren, VA

EXECUTIVE SUMMARY

The *Guide for Incorporating Bioavailability Adjustments into Human Health and Ecological Risk Assessments at U.S. Navy and Marine Corps Facilities, Parts 1 and 2*, has been developed as a resource on assessment of bioavailability for use by Navy Remedial Project Managers (RPMs) and others involved in remediating Navy sites and designing studies to support remediation. The guide brings together the most current information on bioavailability of metals, and synthesizes this information into a practical handbook that explains concepts and identifies types of data that need to be collected to assess bioavailability and incorporate it into risk assessment. Although the guide focuses on bioavailability of metals, many of the basic principles described herein also can be applied to assessing bioavailability of organic compounds.

Part 1: Overview of Metals Bioavailability, contained in this volume, is a primer on the concept of bioavailability and how it can be used in determining risk levels. The *Overview* provides a definition of bioavailability and discusses where bioavailability fits in the risk assessment process for both human health and ecological receptors. This volume provides general information on the types of situations where it may be beneficial to perform the additional studies needed to assess bioavailability and outlines the general factors for determining whether bioavailability studies are appropriate and feasible for a particular site. A brief description of test methods used for assessing bioavailability for human health and ecological risk assessment is provided. The steps in conducting a bioavailability study are outlined and important aspects that affect the acceptability of the results are noted. In addition, a brief summary of metal-specific bioavailability information is presented for those metals that are most often found as contaminants at Navy sites (i.e., arsenic, cadmium, chromium, lead, mercury, and nickel for both human health and ecological risk; and copper, tin and zinc for ecological risk only).

Part 2: Technical Background Document for Assessing Metals Bioavailability, contained in the following volume, provides more in-depth technical information for those professionals involved in designing and performing bioavailability studies. The *Technical Background Document* includes guidelines on the types of studies that need to be performed and methods for collecting data necessary to assess bioavailability with specific considerations for individual metals. Standard operating procedures (SOPs) and suggested protocols for the recommended studies are provided as appendices so that a user can readily access this information.

CONTENTS

ACKNOWLEDGEMENTS	ii
EXECUTIVE SUMMARY	iii
FIGURES	vi
TABLES	vi
ACRONYMS AND ABBREVIATIONS.....	vii
GLOSSARY	ix
1.0 INTRODUCTION.....	1-1
1.1 Why Consider Bioavailability in Risk Assessments?	1-1
1.2 Purpose of the Document	1-2
2.0 WHAT BIOAVAILABILITY IS AND HOW IT IS USED IN RISK ASSESSMENT.....	2-1
2.1 Definitions and Concepts	2-1
2.1.1 Human Health Risk Assessment	2-1
2.1.2 Ecological Risk Assessment	2-2
2.2 Environmental Factors Controlling the Bioavailability of Metals	2-5
2.2.1 Factors Affecting the Mobility of Metals in Terrestrial (Soil) Environments.....	2-5
2.2.2 Factors Affecting the Mobility of Metals in Aquatic (Sediment) Settings	2-7
2.3 How Bioavailability is Incorporated into Risk Assessments	2-7
2.3.1 Human Health Risk Assessment	2-7
2.3.2 Ecological Risk Assessments.....	2-10
3.0 WHEN IT IS APPROPRIATE TO CONDUCT A BIOAVAILABILITY STUDY	3-1
3.1 Where Bioavailability Fits in the Navy’s Tiered Risk Assessment Process	3-1
3.1.1 Human Health Risk Assessment	3-1
3.1.2 Ecological Risk Assessment	3-3
3.2 Situations When Bioavailability Should Be Considered.....	3-5
3.3 General Factors That Determine Whether a Bioavailability Study is Appropriate and Feasible	3-6
4.0 DESIGNING/CONDUCTING A BIOAVAILABILITY STUDY	4-1
4.1 Test Methods for Assessing Bioavailability	4-1
4.1.1 In Vitro Methods for Human Health	4-1
4.1.2 In Vivo Methods for Human Health	4-3
4.1.3 Test Methods for Ecological Receptors	4-5
4.2 Steps in Conducting a Bioavailability Study.....	4-8
4.3 General Considerations	4-8
4.3.1 Human Health Risk Assessments.....	4-8
4.3.2 Ecological Risk Assessments.....	4-12
5.0 CHEMICAL-SPECIFIC CONSIDERATIONS FOR ASSESSING BIOAVAILABILITY TO HUMAN RECEPTORS IN TERRESTRIAL (SOIL) SETTINGS.....	5-1
5.1 Arsenic	5-1
5.1.1 Predominant Forms in Soil.....	5-1
5.1.2 Toxicity Assessment	5-1
5.1.3 Relative Bioavailability Via Oral Exposure.....	5-2
5.1.4 Bioavailability Via Dermal Exposure	5-3
5.1.5 Summary of Pertinent Data.....	5-3

5.2	Cadmium	5-3
5.2.1	Predominant Forms in Soil.....	5-3
5.2.2	Toxicity Assessment	5-3
5.2.3	Relative Bioavailability Via Oral Exposure.....	5-3
5.2.4	Bioavailability Via Dermal Exposure	5-4
5.2.5	Summary of Pertinent Data.....	5-4
5.3	Chromium.....	5-4
5.3.1	Predominant Forms in Soil.....	5-4
5.3.2	Toxicity Assessment	5-4
5.3.3	Relative Bioavailability Via Oral Exposure.....	5-4
5.3.4	Bioavailability Via Dermal Exposure	5-5
5.3.5	Summary of Pertinent Data.....	5-5
5.4	Lead.....	5-5
5.4.1	Predominant Forms in Soil.....	5-5
5.4.2	Toxicity Assessment	5-5
5.4.3	Relative Bioavailability Via Oral Exposure.....	5-5
5.4.4	Bioavailability Via Dermal Exposure	5-6
5.4.5	Summary of Pertinent Data.....	5-6
5.5	Mercury	5-6
5.5.1	Predominant Forms in Soil.....	5-6
5.5.2	Toxicity Assessment	5-7
5.5.3	Relative Bioavailability Via Oral Exposures	5-7
5.5.4	Bioavailability Via Dermal Exposure	5-7
5.5.5	Summary of Pertinent Data.....	5-7
5.6	Nickel.....	5-7
5.6.1	Predominant Forms in Soil.....	5-8
5.6.2	Toxicity Assessment	5-8
5.6.3	Relative Bioavailability Via Oral Exposures	5-8
5.6.4	Bioavailability Via Dermal Exposures.....	5-8
5.6.5	Summary of Pertinent Data.....	5-8
5.7	Relevance to Ecological Receptors in Terrestrial Settings.....	5-8

6.0 CHEMICAL-SPECIFIC CONSIDERATIONS FOR ASSESSING BIOAVAILABILITY TO ECOLOGICAL RECEPTORS IN AQUATIC (SEDIMENT) SETTINGS..... 6-1

6.1	Arsenic	6-2
6.1.1	Predominant Forms in Sediment.....	6-2
6.1.2	Bioavailability and Toxicity in Sediments	6-2
6.2	Cadmium	6-3
6.2.1	Predominant Forms in Sediment.....	6-3
6.2.2	Bioavailability and Toxicity in Sediments	6-3
6.3	Chromium.....	6-4
6.3.1	Predominant Forms in Sediment.....	6-4
6.3.2	Bioavailability and Toxicity in Sediments	6-4
6.4	Copper.....	6-5
6.4.1	Predominant Forms in Sediment.....	6-5
6.4.2	Bioavailability and Toxicity in Sediments	6-5
6.5	Lead.....	6-5
6.5.1	Predominant Forms in Sediment.....	6-5
6.5.2	Bioavailability and Toxicity in Sediments	6-6

6.6 Mercury	6-6
6.6.1 Predominant Forms in Sediment	6-6
6.6.2 Bioavailability and Toxicity in Sediments	6-7
6.7 Nickel	6-8
6.7.1 Predominant Forms in Sediment	6-8
6.7.2 Bioavailability and Toxicity in Sediments	6-8
6.8 Tin	6-9
6.8.1 Predominant Forms in Sediment	6-9
6.8.2 Bioavailability and Toxicity in Sediments	6-9
6.9 Zinc	6-9
6.9.1 Predominant Forms in Sediment	6-9
6.9.2 Bioavailability and Toxicity in Sediments	6-10
7.0 SUMMARY OF SELECTED CASE STUDIES	7-1
8.0 REFERENCES	8-1

ACRONYMS AND ABBREVIATIONS

ABS	absorption fraction
AF	(soil-to-skin) adherence factor
ASTM	American Society for Testing and Materials
AT	averaging time for exposure
ATSDR	Agency for Toxic Substances and Disease Registry
AUC	area under the curve
AVS	acid volatile sulfides
BAF	bioaccumulation factor
BERA	Baseline Ecological Risk Assessment
BJC	Bechtel Jacobs Company, LLC
BRA	Baseline Risk Assessment
BW	body weight
C	concentration
Cal-EPA	California Environmental Protection Agency
CDM	Camp, Dresser, and McKee, Inc.
CEC	cation exchange capacity
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CF	conversion factor
CSF	cancer slope factor
DA	absorbed dose
DAD	dermally absorbed dose
DEQ	Department of Environmental Quality
DTSC	Department of Toxic Substances Control
ED	exposure duration
EF	exposure frequency
Eh	redox potential
EPC	exposure point concentration
ERL	effects range low
ERM	effects range median
EV	(soil contact) event frequency
f_{oc}	fraction organic carbon
GI	gastrointestinal
GLP	Good Laboratory Practice
HCl	hydrochloric acid
HQ	hazard quotient
IR	ingestion rate
IRIS	Integrated Risk Information System
N	normal
NA	not applicable

NEPI	National Environmental Policy Institute
NJDEP	New Jersey Department of Environmental Protection
NOAA	National Oceanic and Atmospheric Administration
OM	organic matter
PBET	Physiologically Based Extraction Test
ppm	parts per million
PRG	preliminary remediation goal
PTI	PTI Environmental Services, Inc.
RAF	relative absorption fraction
RAGS	Risk Assessment Guidance for Superfund
RBC	risk-based concentration
RfD	reference dose
RPM	remedial project manager
SA	(skin) surface area
SAIC	Science Applications International Corporation
SEM	simultaneously extracted metals
SMDP	scientific management decision point
SOP	Standard Operating Procedure
SRA	screening risk assessment
SSSL	site-specific screening level
TBD	to be determined
TCLP	Toxicity Characteristic Leaching Procedure
TOC	total organic carbon
TRV	toxicity reference value
U.S. EPA	U.S. Environmental Protection Agency

GLOSSARY

absolute bioavailability: the fraction or percentage of a compound which is ingested, inhaled, or applied on the skin surface that actually is absorbed and reaches the systemic circulation.

bioavailability: the extent to which a substance can be absorbed by a living organism and can cause an adverse physiological or toxicological response.

cancer slope factor (CSF): the number for a chemical in human health risk assessment used to estimate an upper-bound probability of an individual developing cancer as a result of a lifetime exposure to a particular level of potential carcinogen. Generally, cancer slope factors are available from databases such as U.S. EPA's Integrated Risk Information System (IRIS).

in vivo: within a living organism. In this document, in vivo refers to bioavailability studies conducted using live animals.

in vitro: in an artificial environment outside a living organism. In this document, in vitro refers to bioavailability studies conducted in a laboratory setup that does not use live animals.

reference dose (RfD): the toxicity value for a chemical in human health risk assessment used for evaluating the noncarcinogenic effects that could result from exposures to chemicals of concern. Generally, reference doses are available from databases such as U.S. EPA's Integrated Risk Information System (IRIS).

relative absorption fraction (RAF): the fraction obtained by dividing the absolute bioavailability from soil by the absolute bioavailability from the dosing medium used in the toxicity study from which the reference dose for human health risk assessment was determined.

relative bioavailability: a measure of the extent of absorption among two or more forms of the same chemical (e.g., lead carbonate vs. lead acetate), different vehicles (e.g., food, soil, water), or different doses. In the context of environmental risk assessment, relative bioavailability is the ratio of the absorbed fraction from the exposure medium in the risk assessment (e.g., soil) to the absorbed fraction from the dosing medium used in the critical toxicity study.

toxicity reference value (TRV): an estimate of an "acceptable" chemical dose to a wildlife species used in ecological risk assessment. Toxicity reference values are similar to reference doses used in human health risk assessment but are determined for ecological receptors rather than humans.

1.0 INTRODUCTION

Bioavailability adjustments in risk assessment have recently gained national attention and are becoming increasingly accepted by regulators. Interest in bioavailability is increasing because at some sites consideration of bioavailability has reduced the time and cost necessary for site remediation.

1.1 Why Consider Bioavailability in Risk Assessments?

Bioavailability generally refers to how much of a contaminant is “available” to have an adverse effect on humans or other organisms. Figure 1-1 illustrates the relationship between bioavailability and risk-based cleanup levels. As the figure shows, bioavailability has a direct relationship to exposure dose and risk (i.e., lower bioavailability results in decreased exposure dose and risk). On the other hand, bioavailability is inversely related to risk-based cleanup levels (i.e., lower bioavailability results in increased risk-based cleanup levels). Conversely, higher bioavailability results in increased exposure dose and risk and decreased risk-based cleanup levels. Bioavailability can be influenced by external physical/chemical factors such as the form of a metal in soil or sediment as well as by internal biological factors such as absorption mechanisms within a living organism.




Figure 1-1. Relationship Between Bioavailability and Risk Assessment Endpoints

When risk assessments are adjusted to account for lower site-specific bioavailability, the resulting increase in cleanup levels can in some cases substantially reduce the cost of remediation. A good example is the National Zinc Company National Priorities List (NPL) Site in Bartlesville, OK, where soils and house dust were contaminated with lead, cadmium, and arsenic from smelting activities. The primary concern at this site was the risk to people living in the area, especially children exposed to lead. Remediation to meet the original cleanup goals would have required extensive soil removal and replacement at an estimated cost of \$80 to \$100 million. Determining the site-specific bioavailability was identified as an option for revising the exposure estimates to more realistically reflect the conditions at

this site. The regulators and other stakeholders were consulted from the beginning of the project, a work plan containing detailed protocols for the bioavailability studies was developed, and independent experts were brought in to review the protocols. The bioavailability tests conducted included a rat feeding study to determine the bioavailability of lead and cadmium, and a laboratory extraction test to determine the bioavailability of arsenic. The bioavailability studies indicated that the metals in soil at this site were less bioavailable than had been assumed in the initial risk assessment. By incorporating site-specific bioavailability into the risk assessment, the residential soil cleanup level for lead was increased from 500 mg/kg to 925 mg/kg, the cleanup level for cadmium from 30 mg/kg to 100 mg/kg, and the cleanup level for arsenic from 20 mg/kg to 60 mg/kg, resulting in a reduction in remediation costs for this site of more than \$40 million. In comparison, the cost of planning, conducting, and reporting the bioavailability studies, which took approximately seven months, was approximately \$200,000. Although this example is not typical of the Navy's remediation sites, it does demonstrate how consideration of bioavailability can significantly affect cleanup levels and remediation costs.

1.2 Purpose of the Document

The Guide for Incorporating Bioavailability Adjustments into Human Health and Ecological Risk Assessments at U.S. Navy and Marine Corps Facilities consists of two parts. *Part 1: Overview of Metals Bioavailability*, contained in this volume, is designed for use by remedial project managers (RPMs) and others who want general information on bioavailability. The purpose of the *Overview* is to provide an introduction to the concept of bioavailability (Section 2.0), and to show how it is used in risk assessment and present general guidelines for determining whether bioavailability is worth considering at a particular site (Section 3.0). In addition, the *Overview* provides general information on what a bioavailability study entails and a range of cost, time, and technical requirements needed to conduct such studies (Section 4.0). Profiles of the metals that are most often found to be risk drivers at Navy sites, including arsenic, cadmium, chromium, copper, lead, mercury, nickel, tin, and zinc, are provided in Sections 5.0 and 6.0. These sections briefly summarize for each metal current information that is relevant to determining bioavailability for human health and ecological risk assessment. Finally, a brief review of several case studies is provided in Section 7.0. The scope of this document is limited to bioavailability of metals; however, it should be noted that many of the basic principles described herein also apply to organic compounds.

Part 2: Technical Background Document for Assessing Metals Bioavailability, contained in the following volume, provides more in-depth technical information for those professionals involved in designing and performing bioavailability studies. The *Technical Background Document* includes guidelines on the types of studies that need to be performed and methods for collecting data necessary to assess bioavailability with specific considerations for individual metals. Standard operating procedures (SOPs) and suggested protocols for the recommended studies are provided as appendices so that a user can readily access this information.

2.0 WHAT BIOAVAILABILITY IS AND HOW IT IS USED IN RISK ASSESSMENT

This section defines bioavailability and related concepts, discusses the significant factors that affect the form, distribution, and mobility of metals in soil and sediments, and discusses how quantitative measures of bioavailability can be incorporated into human and ecological risk assessments (Section 4.0 provides a more detailed discussion of how bioavailability is measured).

2.1 Definitions and Concepts

Bioavailability is the extent to which a substance can be absorbed by a living organism and can cause an adverse physiological or toxicological response. For environmental risk assessments involving soil and sediment, this definition implicitly includes the extent to which a substance can desorb, dissolve, or otherwise dissociate from the environmental medium in which it occurs to become available for absorption. For incorporation into a risk assessment, bioavailability must be quantified much like any other parameter in a risk calculation. Thus, it is also useful to define bioavailability in the context of how it is measured.

2.1.1 Human Health Risk Assessment

For human health risk assessment, absolute bioavailability and relative bioavailability are two important and separate measures. ***Absolute bioavailability*** is the fraction or percentage of a compound which is ingested, inhaled, or applied on the skin surface that is actually absorbed and reaches the systemic circulation (Hrudey et al., 1996). Absolute bioavailability can be defined as the ratio of an absorbed dose to an administered dose:

$$\text{Absolute Bioavailability} = \frac{\text{absorbed dose}}{\text{administered dose}} \times 100 \quad (2-1)$$

For studies of absolute bioavailability, the absorbed dose often is determined by measuring the concentration of the compound in blood over time or by measuring the mass of the compound in such excreta as urine, feces, or exhaled air. Internal (i.e., absorbed) doses are useful for characterizing risk if toxicity factors describing the dose-response relationship (i.e., reference dose [RfD], or cancer slope factor [CSF]) are based on an absorbed dose (Figure 2-1). However, because toxicity parameters are generally based on an administered dose rather than an absorbed dose, it is usually not necessary to determine the absolute bioavailability of a contaminant for use in human health risk assessments.

Relative bioavailability is a measure of the extent of absorption among two or more forms of the same chemical (e.g., lead carbonate vs. lead acetate), different vehicles (e.g., food, soil, and/or water), or different doses. Relative bioavailability is important for environmental studies because matrix effects can substantially decrease the bioavailability of a soil- or sediment-bound metal compared to the form of the metal and dosing medium used in the critical toxicity study. In the context of environmental risk assessment, relative bioavailability is the ratio of the absorbed fraction from the exposure medium in the risk assessment (e.g., soil) to the absorbed fraction from the dosing medium used in the critical toxicity study:

$$\text{Relative Bioavailability} = \frac{\text{absorbed fraction from soil}}{\text{absorbed fraction from dosing medium used in toxicity study}} \times 100 \quad (2-2)$$

Relative bioavailability expressed in this manner has been termed the relative absorption fraction (RAF). Incorporation of relative bioavailability (i.e., the RAF) into an exposure assessment results in an improved estimate of the external (i.e., administered) dose (Figure 2-1). It is appropriate to combine the adjusted external dose with toxicity parameters based on an administered dose when characterizing risk.




Figure 2-1. Relationship Between Absolute and Relative Bioavailability and Type of Dose for Risk Assessment

The RAF can be calculated using Equation 2-2 when the absolute bioavailability of a chemical is known for both the dosing medium and the exposure medium. However, as this is almost never the case, a more practical approach is to determine the RAF experimentally with animal (in vivo) studies or laboratory (in vitro) studies without measuring absolute absorption from either the exposure medium or the dosing medium. For example, relative bioavailability can be determined by comparing the fraction of a compound absorbed in a specific target tissue when the compound is administered in soil to the fraction absorbed in the same target tissue when the compound is given in the dosing medium used in the toxicity study.


2.1.2 Ecological Risk Assessment

The uptake by plants and animals of metals from soils, sediments, and water is a complex, dynamic process that involves all levels of the ecological food web. Thus, ecological risk assessment is somewhat more complicated than human health risk assessment. Plants and animals take up bioavailable metals from soils, sediments, and water by contact with external surfaces; ingestion of contaminated soil,

sediment, or water; and inhalation of vapor-phase metals or airborne particles (Brown and Neff, 1993). In addition, animals may take up bioavailable metals from their food. Metal intake may occur through one of these routes of exposure, or through multiple routes functioning either simultaneously or intermittently. A fish, for example, can take up a metal directly from environmental media through its gills, its skin, or through incidental ingestion of sediment; however, it also may ingest and ultimately absorb contaminants through consumption of food (Campbell et al., 1988). Each of these processes involves a different mechanism and, therefore, a different measure of bioavailability.

For ecological evaluations, bioavailability can be addressed using three different approaches (Figure 2-2):

- Evaluating direct exposures to the available fraction of metals present in the environmental media (i.e., sediment or soil)
- Estimating or measuring bioaccumulation directly from the environmental media
- Estimating uptake from ingestion of food.



ECOWEB

Figure 2-2. Illustration of Bioavailability in the Ecological Food Web

Each of these approaches is described below. Because of the complexity of the mechanisms associated with bioavailability in the ecological food web, site-specific factors must be considered prior to incorporating bioavailability adjustments into an ecological risk assessment. Specifically, data evaluated during the planning phase (i.e., problem formulation as defined by the U.S. Environmental Protection Agency [U.S. EPA], 1998g) should be reviewed to determine the relevant exposure pathways and ecological receptors of concern at the site.

Direct Exposures to the Available Fraction in Environmental Media. Metals present in sediments or soils can result in toxicity to organisms directly exposed to them. However, site-specific chemical and physical conditions greatly influence the form in which metals occur in the environment and thus the degree to which they are sorbed to sediments and soils. Therefore, evaluating the total metal concentrations alone does not accurately reflect the fraction biologically available to aquatic and terrestrial organisms. Use of total concentrations as exposure point concentrations (EPCs) in an ecological risk assessment may overestimate actual exposures. Consideration of qualitative and quantitative evidence related to the physical and chemical conditions of a site can assist in determining what portion of the total measured concentration is actually available to organisms exposed. This information provides a better indication of the actual acute and chronic toxicity associated with metals at a site and may help determine which chemicals and/or sampling locations should be included for evaluation in the assessment.

Bioaccumulation from Environmental Media. Another method of evaluating the bioavailability of metals present in soil and sediment is to determine the bioaccumulation of these compounds. This approach provides an estimate of the potential for trophic transfer (i.e., movement of chemicals through the food chain) rather than simply evaluating the potential for direct toxicity to exposed organisms. Bioaccumulation is the uptake and retention of a bioavailable chemical from any one or a combination of possible external sources. Bioavailable metals bioaccumulate by passive diffusion or active transport down a concentration or activity gradient across the outer membranes of the organism (Newman and Jagoe, 1994). As the concentration of the chemical in the tissues increases, the gradient decreases and the rate of loss of the chemical from the tissues tends to increase by either passive diffusion or active transport.

Equilibrium is reached when the rates of uptake and passive or active excretion of the metal are equal. It is necessary to consider bioaccumulation when exposures to upper trophic level species (i.e., birds and mammals) exist.

Uptake from Food. Terrestrial, freshwater, and marine animals are able to accumulate most bioavailable forms of metals from their food. When an animal consumes a lower trophic organism, any metals that have accumulated in the tissues of that organism can be transferred to the consumer (i.e., through trophic transfer). This process occurs primarily or exclusively in the unique environment of the gut of the consumer. Metals that are sorbed or bound to the tissues of a food item and are introduced into the gut of the consumer may be desorbed from the food, dissolved in the gut fluids during digestion, and then partitioned from the gut fluids across the gut lining into the tissues of the consumer. As with uptake directly from soils or sediment, the amount of metal desorbed from the food (i.e., the bioavailable fraction) may be dependent on a number of chemical factors (e.g., chemical form, pH). Consideration of qualitative and quantitative evidence related to the physical and chemical conditions associated with ingestion and absorption can assist in determining what portion of the total measured concentration is actually available to the organisms exposed. This information may help determine which chemicals and/or sampling locations should be included for evaluation in the ecological risk assessment

2.2 Environmental Factors Controlling the Bioavailability of Metals

The bioavailability of an environmental contaminant is largely a function of environmental processes that act on the contaminant to increase or decrease its mobility, thereby making it more or less accessible to the receptor organism. However, physiological factors within the receptor organism, such as acidic gastric juices in the gastrointestinal tract, may also increase the availability of a soil- or sediment-bound contaminant that would otherwise have limited availability under ambient environmental conditions. Thus, for the oral exposure route, there is not an obvious correlation between environmental mobility and bioavailability, so it is important that oral bioavailability studies mimic the physiological conditions under which absorption occurs. For other exposure routes (i.e., dermal absorption, inhalation, and plant uptake), the factors controlling the mobility of the contaminant in the environment also greatly influence the contaminant's bioavailability. Thus, it is relevant to review the processes that affect the fate of a metal in soil and sediment systems.

2.2.1 Factors Affecting the Mobility of Metals in Terrestrial (Soil) Environments

Metals can occur in the soil environment in both the solid phase and the aqueous (i.e., soil solution) phase. In solution, metals can exist either as free ions or as various complexes associated with organic (i.e., functional groups such as carboxyl and phenolic) or inorganic (e.g., anions such as OH^- , CO_3^{2-} , SO_4^{2-} , NO_3^- , or Cl^-) ligands. In the solid phase, metal ions either can be retained on organic and inorganic soil components by various sorption mechanisms (e.g., ion exchange or surface complexation), or can exist as minerals or be co-precipitated with other minerals (e.g., carbonates) in the soil. Ions in solution generally are more available for a variety of processes, including plant uptake and transport; however, metal ions in the solid phase may become available if environmental conditions change.

Dissolution and *precipitation* are the chemical reactions that determine the availability of inorganic *mineral* components of soils. Because most soils are undersaturated with respect to their inorganic mineral components, the minerals undergo continuous dissolution; and, dissolution kinetics is the major factor controlling the availability of mineral-derived metal ions. Some of the more common mineral forms occurring in soils for the metals reviewed in this document are listed in Table 2-1.

Table 2-1. Possible Mineral Species Controlling Soil Solution for Trace Elements (from Hayes and Traina, 1998)

	Aerobic Soils^(a)	Anaerobic Soils^(b)
Arsenic	$\text{Ca}_3(\text{AsO}_4)_2$, $\text{Mg}_3(\text{AsO}_4)_2$, As_2O_5	As, As_2S_3 , As_2O_3
Cadmium	$\text{Cd}(\text{OH})_2$, CdCO_3	Cd, CdS
Chromium	$\text{Cr}(\text{OH})_3$ (low to neutral pH)	$\text{Cr}(\text{OH})_3$
Lead	PbO, PbCO_3 , $\text{Pb}_3(\text{CO}_3)(\text{OH})_2$	Pb, PbS
Mercury	HgCl_2 , HgO, $\text{Hg}(\text{OH})_2$	Hg, HgS
Nickel	NiO, NiCO_3 , $\text{Ni}(\text{OH})_2$	Ni, NiS

(a) Well-drained soils in upland settings (most soils fall into this category).

(b) Seasonally flooded or wetland soils.

The extent to which these mineral species occur in a particular soil and their solubility in various biological fluids (e.g., gastrointestinal tract fluid, sweat, or fluid in the aveoli of the lungs) determines the

relative bioavailability of the various mineral species. In general, the elemental and sulfide forms of a metal are less soluble in biological fluids and hence less bioavailable than the oxide, hydroxide, carbonate, and sulfate forms of the same metal. However, notable exceptions to this rule of thumb exist, such as the following: the elevated pulmonary and dermal bioavailability of elemental mercury; the low solubility of nickel oxides (in the range of nickel sulfide); and the low solubility of chromium hydroxide, the most prevalent form of chromium in soils.

In solution, metals can combine with dissolved organic and inorganic ligands to form complex ions. Examples of such complexes include methylmercury (CH_3Hg^+), cadmium chloride (CdCl^-), and lead bicarbonate (PbHCO_3^+). In general, metals will complex with the most common anions present in soil solution (i.e., inorganic anions such as SO_4^{2-} , NO_3^- , CO_3^{2-} , HCO_3^- , Cl^- , OH^- ; and organic anions such as COO^-). Some metals, such as arsenic and chromium, combine with oxygen to form oxyanions that serve as ligands that can complex with other metals. Arsenite (AsO_3^{3-}), arsenate (AsO_4^{3-}), and chromate (CrO_4^{2-}) are the oxyanions of these metals. The formation of solution complexes can have a significant effect on the mobility of trace metals in soil. For example, trace metals that form chloro-complexes (e.g., CdCl^-) are weakly sorbed and thus likely to be more susceptible to leaching and plant uptake. Although it is likely that different dissolved forms of the same metal will have different absorption efficiencies, it is generally assumed that compounds in the dissolved phase can be completely absorbed regardless of the dissolved species. Therefore, it is generally not necessary to distinguish the dissolved forms of a metal in soil solution for a bioavailability study.

Sorption is an important process because it retains ions on the soil and limits their availability in the soil solution. Sorbed compounds can occur as *surface complexed* (i.e., adsorbed); or, if the density of surface complexes is great enough, as a *surface precipitate* or *cluster* (i.e., a three-dimensional growth on the surface of a soil particle). There is a continuum between surface complexation (adsorption) and surface precipitation such that as the amount of metal coverage increases, surface complexation followed by surface precipitation is the predominant sorption mechanism. The formation of surface complexes (i.e., adsorption) of metals occurs on clay minerals, metal oxides (i.e., hydrous oxides, hydroxides, and oxyhydroxides of iron, manganese, and aluminum), amorphous materials, and organic matter. These soil components contain *surface functional groups* (i.e., molecular units such as hydroxyl, carbonyl, carboxyl, and phenol) that can acquire either a positive or a negative charge, depending on the pH of the soil. Surface complexes can be weakly held (referred to as outer sphere complexes) or more tightly held (referred to as inner sphere complexes) to the soil. Outer sphere complexation is usually a reversible process (i.e., sorption and desorption are identical), whereas inner sphere complexation is often not reversible (i.e., the amount of material desorbed from a soil is less than the amount adsorbed). The non-reversible nature of sorption has been observed for contaminants that have been in contact with the soil for some time, thereby indicating that aged contaminants tend to be less bioavailable than fresh contaminants.

Ion exchange is another type of sorption reaction; however, it is distinguished from the other sorption reactions because it occurs mainly at “fixed charge” sites (i.e., the charge is permanent, not pH dependent) of clay minerals that have undergone isomorphic substitution (i.e., replacement of cations in the clay mineral lattice with other cations of lower charge). Soils with significant negative charge have a high cation exchange capacity (CEC) and low cation mobility. Soils high in clay typically have the highest CEC.

Oxidation-reduction reactions involve the transfer of electrons from one compound to another, resulting in a change in the oxidation state of the compounds involved. The ability of metals to exist in multiple oxidation states is an important property that affects their form and distribution in soils. The most common oxidation states of the soil metals reviewed in this document are as follows: As (III, V), Cd (II), Cr (III, VI), Hg (II), Pb (II), and Ni (II) (copper, tin, and zinc are reviewed in aquatic settings, see Section

2.2.2). Of these metals, only chromium and arsenic are “redox active” (i.e., susceptible to oxidation/reduction reactions) in soil systems. Arsenic exists as As (III) under low redox (i.e., reducing) conditions and as As (V) under high redox (i.e., oxidizing) conditions. Chromium occurs as Cr (III) in most soils under ambient conditions and as Cr (VI) only under highly oxidizing conditions.

In summary, soil conditions that tend to promote precipitation or sorption also tend to reduce the mobility and bioavailability of metals. Thus, the metals that tend to be the most mobile and bioavailable are either those that form weak outer sphere complexes with organic or inorganic (clay, metal oxides) soil components, or those that complex with ligands in solution and are not sorbed. Conversely, metals that form inner-sphere complexes are much less likely to desorb and thus are less mobile and bioavailable. However, in the presence of dissolved organic carbon, the mobility and bioavailability of metals that form inner-sphere complexes may be higher than expected based on sorption behavior, because these metals tend to also form strong soluble complexes. The relative mobility of the metals reviewed in this document is summarized on Table 2-2.

2.2.2 Factors Affecting the Mobility of Metals in Aquatic (Sediment) Settings

Metals are found in all sediments; however, a large amount of the total metals in most sediments is in a residual fraction as part of the natural minerals that make up the sediment particles. These residual metals are not bioavailable. The remaining metals in sediments are adsorbed to or complexed with various sediment components and may be bioavailable (Table 2-3). In oxidized sediments, metals may be adsorbed to clay particles, iron, manganese, and aluminum oxide coatings on clay particles, or dissolved and particulate organic matter. As the concentration of oxygen in sediment decreases, usually due to microbial degradation of organic matter, the metal oxide coatings begin to dissolve, releasing adsorbed metals. In oxygen-deficient sediments, many metals react with sulfide produced by bacteria and fungi to form insoluble metal sulfides. Metals may be released from sorbed or complexed phases into sediment pore water in ionic, bioavailable forms during changes in oxidation/reduction potential. Microbial degradation of organic matter also may release adsorbed metals to pore water. Certain bacteria are able to methylate some metals, such as mercury, arsenic, and lead, to organic species that are more bioavailable than the inorganic forms.

2.3 How Bioavailability is Incorporated into Risk Assessments

It is important to understand how bioavailability data can be used in human health and ecological risk assessments in order to better understand how this parameter should be quantified. Bioavailability is relevant to many aspects of the risk assessment process (e.g., exposure assessment, toxicity assessment); however, this document focuses on the use of bioavailability data to adjust exposure estimates developed in a risk assessment. It should be recognized, however, that other aspects of bioavailability exist that are beyond the scope of this document (e.g., differences in bioavailability between humans and test animals, and variations in the bioavailability of a compound among human subpopulations).

2.3.1 Human Health Risk Assessment

This section illustrates how bioavailability measurements are incorporated into calculations of risk for the oral and dermal exposure pathways, and illustrates how a bioavailability adjustment affects the resulting risk.

For the oral exposure route, relative absorption adjustments can be used to modify the exposure (i.e., intake) estimate (U.S. EPA, 1989). This is illustrated in the following equations, in which the RAF

**Table 2-2. Relative Mobility of Selected Metals in Soil
(from Hayes and Traina, 1998)**

Metal	Most Common Oxidation States in Soil ^(a)	Predominant Forms and Distribution in Soil Systems	Mobility
Arsenic	III	Oxyanion; sorbs more weakly than As(V) to metal oxides and only at higher pH	Moderate
	V	Oxyanion; sorbs strongly to metal oxides; forms relatively insoluble precipitates with iron	Low
Cadmium	II	Cation; sorbs moderately to metal oxides and clays; forms insoluble carbonate and sulfide precipitates	Low to Moderate
Chromium	III	Cation; sorbs strongly to metal oxides and clays; forms insoluble metal oxide precipitates	Low
	VI	Oxyanion; sorbs moderately to metal oxides at low pH, weaker sorption at high pH	Moderate to High
Lead	II (IV)	Cation; sorbs strongly to humus, metal oxides, and clays; forms insoluble metal oxides and sulfides; forms soluble complexes at high pH	Low
Mercury	II (O-I)	Cation; sorbs moderately to metal oxides, and clays at high pH; relatively high hydroxide solubility; forms volatile organic compounds	Low
Nickel	II (III)	Cation; sorbs strongly to humus, metal oxides, and clays; forms insoluble metal oxides and sulfides; forms soluble complexes at high pH	Low

(a) Possible, but less common, oxidation states in soil systems are shown in parentheses; these forms are not discussed.

Table 2-3. Dominant Adsorbed or Complexed Phases of Metals in Oxic and Anoxic Sediments (from Brown and Neff, 1993)

Metal	Associations in Oxic Sediments	Associations in Anoxic Sediments
Arsenic	AsO ₄ ⁻³ -Fe/MnO	As ₂ SO ₃ , AsS, FeAsS
Cadmium	Fe/MnO, OM/S, -CO ₃	CdS
Chromium	OM, FeO	OM, Cr(OH) ₃
Copper	OM, Fe/MnO	Cu ₂ S, CuS, FeCuS
Lead	Fe/MnO	PbS
Mercury	OM	HgS, OM
Nickel	Fe/MnO	OM/NiS, organic thiols
Tin ^(a)	TBT-Cl-OH-CO ₃	TBT-S, OH, -CO ₃
Zinc	Fe/MnO, OM	ZnOM/S

(a) Only butyltins are considered.

CO₃ = carbonates.

FeO = iron oxyhydroxides.

Fe/MnO = iron and manganese oxyhydroxides.

OM = organic matter.

S = sulfides (dominant species given).

TBT-Cl, OH, -CO₃, and -S = tributyltin chloride, hydroxide, carbonate, and sulfide.

expresses the bioavailability of the soil-bound metal compared to the bioavailability of the metal form and dosing medium in the toxicity study from which the CSF or RfD was derived (i.e., $CSF_{administered}$ or $RfD_{administered}$):

$$Risk_{carcinogens} = (Intake \times RAF) \times CSF_{administered} \quad (2-3)$$

$$Risk_{noncarcinogens} = \frac{(Intake \times RAF)}{RfD_{administered}} \quad (2-4)$$

U.S. EPA risk assessment guidance (U.S. EPA, 1989) does not include the RAF term in the risk calculation as shown in the above equation; thus, the U.S. EPA risk equation implicitly assumes a default bioavailability of 1 for the oral pathway. The dermal bioavailability of chemicals in soil is expressed as an absorption fraction (ABS_{soil}) that is incorporated directly into the equation for calculating the dermally-absorbed dose (U.S. EPA, 1992):

$$DAD = \frac{(C_{soil} \times CF \times AF \times ABS_{soil}) \times EF \times ED \times EV \times SA}{BW \times AT} \quad (2-5)$$

where,

- DAD = dermally absorbed dose (mg/kg-d)
- C_{soil} = total concentration in the soil (mg/kg)
- CF = a conversion factor (10^{-6} kg/mg)
- AF = soil-to-skin adherence factor (mg/cm²-event)
- ABS_{soil} = dermal absorption fraction (dimensionless)
- EF = exposure frequency (events/year)
- ED = exposure duration (year)
- EV = soil contact event frequency (events/day)
- SA = skin surface area available for contact (cm²)
- BW = body weight (kg)
- AT = averaging time for exposure (days).

The factors in parentheses describe the absorbed dose per event, DA_{event} (mg/cm²-event). The U.S. EPA (1998f) recommends specific absorption fractions for a few chemicals, and the use of the following default absorption values in the absence of measurements: 1 percent for inorganics and 10 percent for semivolatile organic compounds.

The dermally-absorbed dose is multiplied by the oral RfD or CSF, adjusted to an absorbed-dose basis, to calculate risks via the dermal pathway:

$$Risk_{carcinogens} = DAD \times (CSF_{oral} \times GI_{ABS}) \quad (2-6)$$

and

$$Risk_{noncarcinogens} = \frac{DAD}{(RfD_{oral}/GI_{ABS})} \quad (2-7)$$

Adjustment of the toxicity factors is required because dermal exposures are expressed as an absorbed (i.e., internal) dose, whereas the toxicity factors are usually derived from orally administered doses. GI_{ABS} is the gastrointestinal absorption factor (dimensionless) that expresses the fraction of the orally administered metal in the toxicity study that was absorbed via the GI tract. The U.S. EPA recommends making

adjustments to the toxicity factors only when there is evidence to indicate that the oral absorption in the critical study is significantly less than complete (i.e., <50 percent) (U.S. EPA, 1998g).

2.3.2 Ecological Risk Assessments

As discussed in Section 2.1.2, there are three general approaches that can be used to evaluate bioavailability to ecological receptors. This section explains the methods for including each of these approaches in an ecological assessment.

Direct Exposures to the Available Fraction. In the initial stages of the tiered risk assessment process, estimates of the available fraction of metals in sediment or soil may be limited to a qualitative evaluation of the site-specific chemical and physical parameters that control bioavailability. These data may provide a line-of-evidence argument for inclusion or exclusion of individual chemicals or sampling locations in the risk assessment. The specific parameters considered are discussed further in Section 2.2 and in Sections 5.0 and 6.0 of this document. As the investigation progresses through the tiered evaluation, more complex, quantitative approaches, such as specific analytical techniques or bioassays, may be considered.

For example, as described in Section 4.1.3, analytical techniques may be applied to quantify the specific concentrations of metals in sediments or soils, defined as the simultaneously extracted metals (SEM), that are bioavailable. Concentrations determined from these analytical techniques can be used as adjusted EPCs. For sediments, the estimates of the bioavailable concentration can be further modified based on evaluation of acid volatile sulfides (AVS). In the presence of AVS in sediments, certain metals, including copper, cadmium, lead, nickel, zinc (Ankley, 1996; Ankley et al., 1996), and possibly arsenic and mercury (Luoma, 1989; Allen et al., 1993; Ankley et al., 1996; Neff, 1997a; Berry et al., 1999), precipitate as their respective metal sulfides, which are not bioavailable (DiToro et al., 1990). If the molar concentration of AVS in sediments is higher than the sum of the molar concentrations of these metals in the 1-Normal hydrochloric acid (1-N HCl) extract (the SEM of the sediment), all of the metals are in non-bioavailable forms in the sediments. This relationship can be summarized in the following manner:

SEM:AVS > 1, metals are present in bioavailable forms

SEM:AVS < 1, metals are not likely to be bioavailable.

If the SEM:SVS>1, then these data can be used to calculate an EPC as discussed below. It is important to note that each of the metals evaluated has a different binding affinity for sulfides (U.S. EPA, 1994a). Currently there is considerable debate regarding the relative affinities of each of the metals; however, typically it is assumed that at equilibrium, copper will preferentially react with AVS, displacing all other metals. If the available AVS is not completely saturated by copper, then the remaining metals will react in the following order: lead, cadmium, zinc, and nickel. In this model, the amount of copper in the sediment that is potentially bioavailable and toxic is considered to be defined as follows:

$$Cu_b = (Cu_{SEM} - AVS) * (MW_{cu}) \quad (2-8)$$

where,

Cu_b = concentration of copper that is bioavailable (mg/kg)

Cu_{SEM} = molar concentration of Cu as defined by simultaneous extraction (moles/kg)

AVS = molar concentration of AVS (moles/kg)

MW_{cu} = molecular weight of copper (mg/moles).

The bioavailable concentration of the other metals in sediment may be determined in the same manner, following the order described above. For each successive metal, the molar concentration of AVS applied should be decreased according to the molar concentration of the preceding chemical; when the concentration of AVS is zero, all remaining metals are assumed to be bioavailable. The metal concentrations derived in this manner can be used as EPCs.

Bioaccumulation from Environmental Media. Uptake of sediment-bound or soil-bound metals by organisms (i.e., bioaccumulation) either may be measured directly by collecting and analyzing the tissues of representative organisms, or may be estimated (BJC, 1998). In the initial stages of a risk assessment, estimates are typically derived according to the following equation:

$$C_t = C_s * BAF \quad (2-9)$$

where,

C_t = concentration in tissue (mg/kg)

C_s = concentration in sediment or soil (mg/kg)

BAF = bioaccumulation factor ($[mg/kg_{tissue}] / [mg/kg_{sed/soil}]$).

In the event that tissue-based TRVs are available, C_t can be used to derive a hazard quotient (HQ) as defined by the equation:

$$HQ = \frac{C_t}{TRV} \quad (2-10)$$

In addition C_t can be used to represent the exposure point concentration for estimating ingested doses for upper trophic level species. For example:

$$Dose_{Ingested} = \frac{C_t * IR}{BW} \quad (2-11)$$

where,

IR = ingestion rate of receptor species (kg/day)

BW = Body weight of receptor species (kg).

BAF values, defined as the ratios of the concentration of the chemical in the tissues of the organism to the concentration of the chemical in sediment or soil, have been derived for various chemicals and species and are available in the literature. In the event that BAF values for relevant chemicals or species are not available in the literature, they may be derived using tissue and soil or sediment data available in the literature or determined experimentally at the site. This relationship may not be valid for those metals that are essential trace nutrients for plants and animals.

Uptake from Food. For upper trophic level species, quantitative data also can be used to modify ingested doses for use in calculating risk estimates. These data would be incorporated as described for the noncarcinogenic human health risk assessment. For example, when evaluating exposures resulting from the ingestion of contaminated prey items, the following simplified equation may be used to determine the risk from food ingested by the ecological receptor:

$$Risk = (Intake \times ABS) / TRV \quad (2-12)$$

where,

Intake = ingested dose (mg/kg/day)

ABS = absorption factor (unitless)

TRV = toxicity reference value (mg/kg/day).

For screening-level evaluations, the ABS is typically assumed to be 1 (i.e., absorption is 100 percent). However, as the investigation progresses through the ecological risk assessment process, it may be possible to refine this value to reflect actual conditions either through a review of the relevant literature, or through bioassays as described for human health exposures.

3.0 WHEN IT IS APPROPRIATE TO CONDUCT A BIOAVAILABILITY STUDY

This section discusses a variety of considerations that RPMs should review when deciding if a bioavailability study makes sense for their site. Section 3.1 discusses where in both the human health and the ecological risk assessment processes it is appropriate to conduct a bioavailability study. Section 3.2 outlines several situations where bioavailability might offer an appropriate solution to a given remediation problem, and Section 3.3 discusses factors that may affect whether a bioavailability study is worthwhile for a particular site.

3.1 Where Bioavailability Fits in the Navy's Tiered Risk Assessment Process

The Navy has applied tiers to the risk assessment process for assessing human and ecological risks (see Figures 3-1 and 3-2). This section briefly discusses the major steps in the tiered risk-assessment process and where it is appropriate to conduct a study to support a site-specific bioavailability adjustment.

3.1.1 Human Health Risk Assessment

Figure 3-1 illustrates the Navy's three-tiered *human health risk assessment process*. Bioavailability data can be incorporated during the risk-based screening step (Tier I) and during the Baseline Risk Assessment (BRA) (Tier II) because both steps rely on the use of exposure and risk calculations that allow for the incorporation of bioavailability adjustments. Tier I involves a risk-based screening step in which site concentrations are compared to generic or site-specific risk-based screening levels. Sources of generic screening levels include the U.S. EPA Region III risk-based concentrations (RBCs) (U.S. EPA, 2000) and the U.S. EPA Region IX preliminary remediation goals (PRGs) (U.S. EPA, 1999). Another source of generic screening levels for soil is Appendix A of the *Soil Screening Guidance: Technical Background Document* (U.S. EPA, 1996a). The Region III RBCs and Region IX PRGs are updated periodically as new toxicity and physio-chemical data become available, whereas the values in the *Soil Screening Guidance* have not been updated since the document was issued. Therefore, stakeholders need to decide which screening values to use for a particular site. Bioavailability data are not incorporated into the generic Tier I screening values because the Tier I values are based on conservative default exposure assumptions designed to provide screening levels protective of most sites across the country.

If site concentrations exceed the generic Tier I values, site-specific screening levels (SSSLs) are calculated in Tier IB and compared to site concentrations (Figure 3-1). SSSLs differ from the generic Tier I screening levels in that actual physical properties of the site are incorporated into the SSSL calculations in place of default values inherent in the generic "look-up" values. In addition, whereas generic Tier I screening levels are available for only specific exposure scenarios (typically ingestion, dermal contact, inhalation of vapors and particulates), SSSLs can be developed for other relevant pathways (e.g., food ingestion, vapor intrusion to buildings) or to take into account indirect exposure scenarios (i.e., when receptors are exposed to contaminants that are transported from the source to other exposure media such as groundwater or air). Because the Tier I SSSLs are calculated values rather than "look-up" values, Tier IB provides an opportunity for the incorporation of bioavailability data. Several resources are available for developing SSSLs, including Part B of the U.S. EPA's *Risk Assessment Guidance for Superfund (RAGS)* document (U.S. EPA, 1991a), the *Soil Screening Guidance: Technical Background Document* (U.S. EPA, 1996a), and the American Society for Testing and Materials *Standard Guide for Risk-Based Corrective Action Applied at Petroleum Release Sites* (ASTM, 1995) and *Standard Provisional Guide for Risk-Based Corrective Action* (ASTM, 1998).




Figure 3-1. Incorporating Bioavailability in the Tiered Human Health Risk Assessment Process

Under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), the risk-based screening step (i.e., Tier I) allows areas of the site with contaminant concentrations below the risk-based screening levels to be eliminated from further action; whereas, areas of the site with contaminant concentrations above the soil screening levels must undergo further assessment (U.S. EPA, 1994a, 1994b, and 1996a). Further assessment may involve conducting a BRA, although site owners can elect to bypass the BRA and remediate the site to the soil screening levels. Because Tier I provides a means for eliminating low-risk sites early in the CERCLA process, consideration should be given to conducting a bioavailability study (in Tier IB) to support the calculation of realistic risk-based screening levels.


Tier II of the human health risk assessment process involves conducting the BRA (Figure 3-1). The U.S. EPA's RAGS document (U.S. EPA, 1991a) provides guidance on conducting a human health BRA. A BRA involves four basic steps: data collection and evaluation, exposure assessment, toxicity assessment, and risk characterization. As discussed in Section 2.3.1, bioavailability data can be incorporated in the BRA to adjust exposure estimates for key pathways (e.g., oral), or to extrapolate toxicity data from one route of exposure to another (e.g., GI absorption data are required to adjust oral toxicity factors to an absorbed-dose basis for calculating dermal risks). If bioavailability data are to be incorporated into the BRA, a site-specific bioavailability study is needed early in the BRA to provide the necessary data for making these adjustments. The results of the Tier I assessment can provide an early indication as to whether or not a bioavailability study might be necessary during the BRA, as this information is useful for identifying contaminants and exposure routes that present the highest risks for the site.

Tier III of the human health risk assessment process involves an assessment of the risks associated with various remedial alternatives. Guidance for evaluating short-term and long-term risks associated with site remediation activities is provided in Part C of the U.S. EPA's RAGS document (U.S. EPA, 1991b). If these risks are assessed in a quantitative manner, incorporation of bioavailability data may be appropriate in this phase of the risk assessment process.

3.1.2 Ecological Risk Assessment

Figure 3-2 illustrates the incorporation of the bioavailability evaluation into the Navy's *ecological risk assessment* process. As indicated, the first step in an ecological risk assessment (Tier 1) is a Screening Risk Assessment (SRA). This step is a conservative, worst-case evaluation of the potential risks at the site. Therefore, all chemicals are assumed to be 100 percent bioavailable. All pathways are identified, and EPCs are determined for all relevant environmental media. Toxicity benchmarks are identified based on available water, sediment, and soil criteria. If the EPCs do not exceed the selected toxicity benchmarks, the site passes the SRA and is closed out for ecological concerns. If the EPCs exceed the selected toxicity benchmarks, the site either has an interim cleanup, or proceeds to the second tier.

Tier 2, the Baseline Ecological Risk Assessment (BERA), entails a more detailed, less conservative approach incorporating site-specific exposure factors. Bioavailability considerations may be incorporated into this tier as part of Step 3a (Refinement of Conservative Exposure Assumptions) in a number of ways, depending on the data, funding, and time available. For example, as a first effort, chemical and physical parameters, such as sediment and soil pH, total organic carbon (TOC), redox potential (Eh), specific form of the metal, SEM/AVS, can be evaluated. Evaluation of each of these factors provides qualitative information for use in a line-of-evidence approach to eliminating individual metals or the site from future consideration. Similarly, application of literature-based bioaccumulation factors or absorption fractions, if appropriate, can provide evidence demonstrating a lack of bioavailability. If, based on these refinements, evidence indicates that the site poses acceptable risks, then the site exits the ecological risk



Note: Modified from the Navy Ecological Risk Assessment Tiered Approach (<http://web.ead.anl.gov/ecorisk>), which is based on the U.S. EPA's 8-Step Ecological Risk Assessment Process.

Figure 3-2. Incorporating Bioavailability in the Tiered Ecological Risk Assessment Process

assessment process. Otherwise, the evaluation proceeds to Step 3b, which involves a more extensive evaluation of site-specific information.

In Step 3b, additional site-specific data may be collected, such as concentrations of metals in tissues of organisms from the site, or measurement of the bioavailable fraction in sediment or soil through sequential extraction techniques. In addition, site-specific bioassays such as bioaccumulation tests or relative bioavailability are considered. It is important to note that site-specific information collected previously should be carefully evaluated to determine the potential cost-effectiveness of proceeding with these more expensive and time-consuming bioassays. If determined to be appropriate, the results of these tests, combined with the data previously collected, can be evaluated to determine if the site poses acceptable risks. If the risks are determined to be acceptable, no further evaluation or remediation from an ecological perspective is required. If the risks are determined to be unacceptable, and additional evaluation in the form of remedy development is appropriate, the process proceeds to the third tier.

The focus of the Tier 3, Evaluation of Remedial Alternatives is to develop site-specific, risk-based cleanup goals and to determine the appropriate remedial strategy. All site information collected during the assessment, including that pertaining to the potential for bioavailability, should be evaluated when considering the various remedial alternatives.

3.2 Situations When Bioavailability Should Be Considered

Several types of situations where bioavailability studies might be beneficial are described below. Note, however, that there can be many other site-specific factors or conditions that ultimately determine whether bioavailability studies are worth pursuing for a given site (see Section 3.3).

- **When a risk estimate slightly exceeds an acceptable level and triggers a requirement for remediation.** If it can be shown that the contaminant at the site is less available to the receptor than was assumed in the initial risk assessment, the risk estimate potentially could be reduced below the acceptable limits, thus avoiding remediation while still being protective of human health and environment.
- **When risk-based cleanup goals require extensive and/or expensive remediation.** This situation includes sites with large areas of elevated contaminant concentrations over much of the site as well as sites where remediation to reach the required cleanup goal is very expensive. In these cases, if it can be demonstrated that the contaminant at the site is less available than was assumed in the original risk assessment, the risk-based cleanup goals can be higher. Higher cleanup goals potentially could reduce the area or volume of soil that requires remediation or increase the concentration that must be achieved by remediation. At the Butte, MT Superfund site where mining activities had resulted in widespread lead contamination, bioavailability studies found that availability of lead from soil at the site was only 12 percent compared to the default assumption of 30 percent. As a result, the cleanup goal for lead was increased from the default of 500 ppm to 1,200 ppm, and tens of millions of dollars were saved in cleanup costs.
- **When remediation is not technically feasible.** In this case, either the required remediation cannot be carried out due to site conditions or an effective remediation technology does not exist to achieve the required cleanup goals. If the contaminants at the site are less bioavailable than was assumed in the initial risk assessment, the risk estimate might be decreased to an acceptable level or calculation of risk-based cleanup goals might yield higher goals that are feasible to achieve.

- **When remediation activities will adversely impact the environment.** In some cases, the remediation activities required to achieve the cleanup goals for a site would have adverse impacts on the environment. Such impacts include habitat destruction, increased potential for erosion, or re-release of contaminants into other environmental media. At the East Fork Poplar Creek site in Tennessee, mercury contamination was spread over 650 acres of the creek's forested watershed. Further study revealed that most of the mercury was in a form that has low bioavailability. This was confirmed by animal uptake and simulated human digestion studies. Cleanup goals were adjusted from the original goal of 10 ppm, based on methylmercury, to 400 ppm. Cleanup costs were cut from an estimated \$1.2 billion to approximately \$8 million, while leaving a large tract of wildlife habitat undisturbed (NEPI, 1998).

3.3 General Factors That Determine Whether a Bioavailability Study is Appropriate and Feasible

This section highlights general factors that an RPM should consider in deciding whether a site-specific bioavailability study is likely to be beneficial for a site.

- **Number of chemicals driving risk.** If three or fewer chemicals drive the risk at a site, then it is possible that bioavailability adjustments could reduce risk estimates enough to justify the cost of doing the bioavailability study. If more than three chemicals drive the risk, a bioavailability adjustment of only a few may not decrease the risk estimate sufficiently to justify the cost of the study.
- **Form of the chemical or the exposure medium for the site compared to the reference dose.** If the form of the chemical found at a site is different than the form used in the toxicity study on which the reference dose is based, then the bioavailability of that compound may be different and conducting a site-specific bioavailability study potentially could result in a significant reduction in risk. An example of this situation is when the form of metal used in a toxicity study is a very soluble form (as is often the case), and the form of metal found in soil has a low solubility. Also, if the exposure medium is different between the reference dose toxicity study and the site (e.g., reference dose was given in water while site exposure is to soil), the bioavailability at the site may be sufficiently different from that reported in the toxicity study to justify a bioavailability study. If the forms or exposure media are similar, then bioavailability is more likely to be similar and a bioavailability adjustment may not be worthwhile.
- **Potential for regulatory acceptance.** Although most regulatory policies allow for bioavailability adjustments, there is no requirement that these adjustments be considered or accepted by the regulators. Therefore, it is important to consider the regulatory climate for the site before undertaking a bioavailability study. The regulators for the site should be contacted to determine if they are receptive to the concept of a bioavailability adjustment. Also, it may be helpful to determine whether there are any precedents for approval of bioavailability adjustments by that agency.
- **Whether bioavailability studies can be completed within the required time frame for the site.** The time required for a bioavailability study can vary depending on the type of study required to collect the necessary data. Generally, simple in vitro (laboratory) tests require less time than in vivo (live animal) feeding studies. More detailed information on time required for various types of studies is provided in Section 4.3.

- **The cost of bioavailability testing compared to the cost of cleanup.** The cost of performing bioavailability studies and incorporating the results into risk assessment must be weighed against the cost of cleanup and the potential cost savings that could result from the bioavailability study. Costs of bioavailability studies can vary substantially depending on what tests are done and who is selected to do them. Section 4.3 provides some rough guidelines on the costs of various types of studies.
- **Existing site data support a bioavailability study.** Information commonly collected during a site investigation should be reviewed when evaluating whether to proceed with a site-specific bioavailability study. Both historical site information and soil parameter data bear on the likely results of such a study. Under certain circumstances, it may be possible to use existing site data to indicate the likely outcome of a bioavailability study, and thereby help determine whether to proceed with the study itself. In general, however, site data cannot be used in place of site-specific bioavailability studies. The following information on using site data to “estimate” bioavailability is intended as a general guideline; soils at specific sites may not conform to all of the general trends discussed here. Furthermore, the generalizations apply mainly to the oral (ingestion) exposure route, which has been the most extensively studied to date. The impact of site history and soil chemistry parameters on the oral bioavailability of metals from soil is indicated in Table 3-1.
 - Historical site information to consider includes both the types of metals contamination present and the length of time that the contamination has been resident in soils or sediments (i.e., the weathering or aging time). The source of contamination can indicate the likely forms in which the metals were deposited in the soils. In general, soils that contain sulfide or elemental metal forms yield lower bioavailability values than soils that contain oxide or carbonate metal forms. Nickel is a notable exception to this trend, and forms several insoluble oxide species. In addition, small mineral particles yield higher bioavailability than large mineral particles. Soil weathering reactions change the bioavailability of metals over time. In general, metal forms with high bioavailability (oxides and carbonates) alter to less bioavailable forms, while metals with low bioavailability (sulfides and elemental forms) alter to more bioavailable forms. The length of time that the metals have been present in the soil will determine the extent of these weathering reactions, and the current bioavailability of the metals in soil.
 - Site-specific soil chemistry determines the products of the soil weathering reactions discussed above. Measurements of soil parameters such as pH, TOC, total carbonate (alkalinity), and iron and manganese concentrations may therefore indicate the likely outcome of a site-specific bioavailability study. In general, weathering products that form in acidic soils (pH less than 5.0) are more stable, and less bioavailable, in the acidic environment of the stomach, while weathering products from alkaline soil environments (pH greater than 8.0) yield elevated bioavailability values.
 - Most of the metals reviewed in this document (cadmium, lead, mercury, and nickel) can alter to carbonate forms in alkaline soils, and these carbonate metal forms are highly bioavailable via the oral exposure route. Soils containing elevated TOC (greater than 5 to 10 percent) tend to contain metals that are complexed to organic matter; these organically complexed metals appear to have elevated oral bioavailability (this is particularly true for lead and mercury). These same soils/sediments will often contain relatively insoluble sulfides as a result of the action of sulfate-reducing bacteria. This mechanism is limited to cadmium, mercury, lead, and nickel in seasonally flooded soils. Finally, soils with elevated iron and manganese

concentrations (greater than 3 to 5 percent combined) tend to have reduced bioavailability, particularly for arsenic due to increased sorption on these soil components.

- The research to date indicates that regulatory leaching tests, such as the Toxicity Characteristic Leaching Procedure (TCLP), do not predict the oral bioavailability of metals from soil. Therefore, results from TCLP testing should not be used in estimating the extent of metals bioavailability from soil.

Table 3-1. Impact of Site History and Soil Chemistry on the Oral Bioavailability of Metals

Site History	Bioavailability		
	Low	Medium	High
Metal Forms:			
Sulfides	X		
Elemental (metallic)	X		
Sulfates		X	
Carbonates			X
Oxides			X (except Ni)
Particle Size (of metal-bearing grains):			
Small			X
Large	X		
Weathering/Aging Time:			
Sulfides	X	→	
Elemental	X	→	
Carbonates			X ←
Oxides			X ←
Soil Chemistry			
pH:			
Acidic		X	
Basic			X (Cd, Hg, Pb, Ni)
Alkaline soils			X (Cd, Hg, Pb, Ni)
High TOC			X (Hg, Pb)
High Fe and Mn		X (As)	
Sulfide-producing soil		X (Cd, Hg, Pb, Ni)	

4.0 DESIGNING/CONDUCTING A BIOAVAILABILITY STUDY

For assessing potential human health risks, bioavailability adjustments usually must be supported by a site-specific study because it generally is not possible to predict the bioavailability of a compound based on other, more fundamental physical or chemical properties of the site or the contaminant. For ecological risk assessments, there are a variety of ways to incorporate bioavailability, and adjustments can be determined either experimentally or with estimation techniques (e.g., bioaccumulation is often modeled using literature-derived bioaccumulation factors). This section provides background information on the types of tests that can be used to assess the bioavailability of a metal to human and ecological receptors and the resources (i.e., cost, time, and technical expertise) required to conduct such tests. The discussion is presented from the perspective that a site-specific bioavailability study will be designed and conducted during risk assessment activities. Thus, recommendations are offered regarding the appropriate steps to include in a bioavailability study to ensure that the study is acceptable to involved regulatory agencies.

4.1 Test Methods for Assessing Bioavailability

A wide variety of methods have been used to study the bioavailability of metals in soils and sediments. For soils, the focus has been on studies in laboratory animals and simple in vitro extraction tests to assess the oral bioavailability of metals in soils relative to the bioavailability of more soluble metal compounds. Most of these studies have been conducted for use in human health risk assessment. For sediments, the bioavailability of metals to ecological receptors has been the focus of most research to date.

For all of these studies, a critical finding is that site-specific studies are generally required. Studies conducted using soluble metal compounds freshly mixed with soil or sediment generally do not show significant reductions in bioavailability, and will not provide a representative indication of the relative bioavailability of metals in soil or sediment at a specific site. Consequently, studies must be conducted using weathered soils or sediments. In addition, it is important that the samples being tested be characterized for parameters such as pH, TOC, CEC, particle size (sand, silt, clay), total metals (Fe, Mn, Al), and available anions (PO_4 , SO_4 , CO_3). Also, it is also important that, for studies predicting human oral absorption of metals in soils, the soils be sieved to include particle sizes of less than 250 microns, because it is these finer particles that are thought to adhere to hands and be ingested during hand-to-mouth activities. For dermal absorption studies, particle sizes of less than 150 microns are the most likely to adhere to skin.

4.1.1 In Vitro Methods for Human Health

This section describes the application of simple laboratory extraction tests (in vitro tests) that are predictive of the bioavailability of metals from soil to humans. These methods are both rapid and inexpensive, requiring only a day to conduct and costing only a small fraction of what an in vivo study (discussed below) would cost. Although in vitro work has focused primarily on determining the oral bioavailability of arsenic and lead, results from these two elements can be extrapolated to other metals based on universal solubility-limiting factors and similarities in the aqueous geochemistry of certain elements. In addition, the dermal absorption of chromium from soil and waste materials has been evaluated by extraction tests using both real and synthetic human sweat (Horowitz and Finley, 1993; Wainman et al., 1994).

Simple extraction tests have been used for several years to assess the degree of metals dissolution in a simulated GI-tract environment (Ruby et al., 1993, 1996, and 1999). The predecessor of these systems was developed originally for nutrition studies to assess the bioavailability of iron from food (Miller et al., 1981; Miller and Schricker, 1982). In these systems, various metal salts, or soils containing metals, are

incubated in a low-pH solution for a period intended to mimic residence time in the stomach. The pH then is increased to near neutral, and incubation continues for a period intended to mimic residence time in the small intestine. Enzymes and organic acids are added to simulate gastric and small-intestinal fluids. The fraction of a metal that dissolves during the stomach and small-intestinal incubations represents the fraction that is bioaccessible (i.e., is soluble and available for absorption).

The currently available in vitro tests (Medlin, 1997; Rodriguez et al., 1999; Ruby et al., 1996) are designed around human pediatric gastrointestinal conditions, and are intended to mimic fasting conditions. Critical design factors that have been evaluated include extraction fluid chemistry and temperature, extraction time, mixing rate, and the particle size of the test material. Because the goal is to develop the simplest test possible, which will yield the highest repeatability and reproducibility, these tests have been streamlined to include only those factors that control the dissolution of a particular metal.

The research to date indicates that the fractional extraction of arsenic or lead during a one-hour incubation in acidic fluid (pH 1.5 in hydrochloric acid) is a good surrogate for relative arsenic or lead bioavailability values derived from in vivo studies (Medlin, 1997; Rodriguez et al., 1999; Ruby et al., 1996). Figure 4-1 shows the correlation of in vivo and in vitro tests for lead bioavailability. Most laboratories currently are using a specialized test cell (Figure 4-2) for these studies; however, Rodriguez et al. (1999) replaced this cell with mason jars and achieved equally good results. It is important to maintain a constant pH during the test (i.e., 1.5 ± 0.3), because the solubility of most metals is highly pH dependent, and allowing the pH to fluctuate may influence the test results. Note that incorporating the food material used during the Rodriguez et al. (1999) studies of arsenic bioaccessibility is not recommended, because the food material contained elevated phosphate concentrations (nearly 3 percent available phosphate), which enhanced the solubilization of soil arsenic.




Figure 4-1. In Vitro to In Vivo Correlation for Lead in Soil

No published in-vitro-to-in-vivo correlations exist for cadmium, chromium, mercury, or nickel. Because all of these metals may occur in soil as discrete mineral forms with varying oral bioavailabilities, it appears that the same controls on bioavailability will be in effect for these metals as those for arsenic and




Figure 4-2. In Vitro Test System

lead. At this time, it is recommended that the in vitro test, which consists of a stomach-phase (i.e., acidic) incubation, be applied to determining the bioaccessibility of arsenic, cadmium, lead, and nickel from soil. Chromium and mercury are best evaluated using sequential stomach-phase and intestinal-phase incubations.

Before undertaking an in vitro study, it is important to consider the desired use for the data. Will the data be used primarily as a range-finding tool, and for guiding further study of site soils using an in vivo model, or are the data intended for use in making a quantitative adjustment to a human health risk assessment? If it is the latter, it is critical to establish a dialogue with the relevant regulatory agency as early as possible, because the use of in vitro data for making adjustments to human health risk assessments is not widely accepted by regulatory toxicologists. Submittal of a study protocol to the regulatory agency is generally a good place to start the dialogue over study design issues and the acceptable uses for these types of data. Appropriate protocols (i.e., Standard Operating Procedures [SOPs]) for in vitro methods may be found in Part 2 of this Guide.

4.1.2 In Vivo Methods for Human Health

Most of the in vivo research to date has focused on the oral bioavailability of metals in soils. This focus reflects the observation that human health risk-based soil cleanup levels for metals are typically driven by ingestion exposures. New dermal exposure guidance from U.S. EPA (1998f) that includes default assumptions of 1 percent dermal bioavailability for most metals (3 percent for arsenic) will cause dermal exposures to be important at some sites in the future. Consequently, this section focuses on methods for assessing oral bioavailability using laboratory animals. Dermal absorption studies are described briefly. Inhalation studies are not discussed because site-specific studies will seldom be relevant, as inhalation is

not a pathway that contributes significantly to risk from metals in soil. When evaluating whether to conduct a bioavailability study, and what form it should take, the Data Quality Objectives (U.S. EPA, 1994b) process should be used to develop the study.

Although the oral bioavailability study methods described are generally used for studies in laboratory animals, it is useful to note that many of these same methods may be used for studies in humans. Recently, lead bioavailability studies in humans have been conducted. The protocols for these studies must undergo scrutiny by institutional review boards to ensure that no unacceptable risks will be imposed, and that informed consent will be obtained.

Oral bioavailability studies generally involve measuring chemical concentrations in body tissues or excreta at various time points after dosing. The specific study design needs to be selected after considering how the metal being studied is handled by the body. Some metals are well absorbed and rapidly excreted in the urine (arsenic is a good example), while other chemicals may have more limited absorption and may accumulate in body tissues. For example, lead is accumulated in bone, while cadmium is accumulated in the kidneys and liver. Different study designs are needed to reflect these different characteristics. Thus, there is no one oral bioavailability study protocol that can be applied uniformly to all metals.

The four primary methods used to study the oral bioavailability of metals are:

- **Measurement of blood concentrations over time for oral and intravenous doses.** The area under the curve (AUC) is calculated, and oral absorption is determined by comparing the AUC_{oral} to the $AUC_{\text{intravenous}}$ (see Figure 4-3). This method works best for metals that are well absorbed, and rapidly and completely excreted (e.g., arsenic).




Figure 4-3. Comparison of AUCs for Blood Concentrations

- **Measurement of the fraction of the dose that is excreted in the feces.** This measurement generally reflects unabsorbed metal, so absorbed dose is calculated by subtracting the excreted dose from the administered dose. This method may underestimate absorption if a metal is absorbed, then excreted via bile back to the gastrointestinal tract.
- **Measurement of the fraction of the dose that is excreted in urine.** This fraction provides an estimate of absorbed dose for metals that are rapidly excreted primarily in the urine (e.g., arsenic).
- **Comparison of tissue concentrations after administration of different forms of a metal.** This method provides an estimate of relative bioavailability, and is most useful for metals that are preferentially accumulated in specific tissues.

For all of these methods, if metals in soil are compared to a soluble form of the metal, the resulting relative bioavailability estimate may be used to derive exposure estimates. The specific animal model selected for use in the studies should be based on an understanding of the behavior of the metal being studied in that animal, and on any significant differences between the animal selected and humans. Other factors to consider include the age of the animals (for example, lead is absorbed more completely in young animals), and the nutritional status and diet for the animals (for example, lead is better absorbed in fasted animals).

A study protocol or work plan must be prepared that specifies dose levels, frequency of dosing, number of animals per group, samples to be collected and the timing and frequency of sample collection, and quality assurance procedures to be followed. The U.S. EPA has issued specific regulations for quality assurance for laboratory studies called Good Laboratory Practices (GLPs) (40 CFR Part 792). These regulations specify the elements to be included in a study protocol, and quality assurance procedures to follow. It is advisable to require a contractor to conduct studies in accordance with the GLPs.

The preferred methods for studying dermal absorption of metals include in vivo studies and in vitro studies. Rhesus monkey and swine are useful animal models for in vivo dermal studies. In vitro dermal studies are performed using human cadaver skin. No simple in vitro extraction methods have been developed for routine use in screening a series of site soils for relative dermal bioavailability. In designing dermal absorption studies for use in risk assessment, it is critical that the nature of potential exposures be mimicked as closely as possible. Critical factors include the use of a fine fraction of the soil (particles less than 150 microns are thought to be most likely to adhere to skin), the use of a soil load that will not exceed a monolayer on the skin surface (generally less than 5 mg soil/cm² of skin), and an exposure period representative of expected exposures at the site. An extensive review of methods for studying dermal absorption can be found in the U.S. EPA's *Dermal Exposure Assessment* document (1992).

4.1.3 Test Methods for Ecological Receptors

As discussed in Sections 2.1.2 and 2.3.2, a variety of approaches may be used to incorporate bioavailability into ecological risk assessments. For each of these approaches, several specific test methods may be used to provide a quantitative or qualitative measure of the bioavailable metals depending on the complexity of the site and the current phase of the risk assessment process (i.e., Tier 1 or Tier 2). In general, the more qualitative methods are typically used in the initial stages of the Baseline Ecological Risk Assessment while the site-specific bioassays or complex analytical techniques are reserved for consideration as the risk assessment process progresses (Figure 3-2). Table 4-1 summarizes the test methods associated with each of the approaches discussed.

Table 4-1. Test Methods for Assessing Bioavailability in Ecological Risk Assessments

Approach	Methodology	Purpose	Limitation
Direct Exposures to the Available Fraction in the Media	Extraction Techniques (e.g., 1-N HCl)	Provides numerical estimate of bioavailable fraction (i.e., concentration)	No single extraction technique has been demonstrated to completely characterize the bioavailable fraction
When evaluating direct exposures/toxicity associated with sediments or soils	Comparison of AVS/SEM (sediment only)	Provides additional modification to bioavailable fraction estimate	Recent data indicate that the AVS/SEM model is not always a good predictor
	Evaluation of chemical and physical parameters	Provides qualitative evidence for line-of-evidence argument	Evidence is only qualitative
	Toxicity tests	Absence of toxicity provides line-of-evidence support for lack of bioavailability	Results of toxicity tests can be difficult to interpret and may be costly and time consuming
Bioaccumulation from Environmental Media	Collect and analyze site specific tissue data	Provides a measure of amount of chemical that is taken up by resident species	Measured concentrations may be impacted by sources other than those at the site
When estimating tissue concentrations to evaluate trophic transfers	Estimate tissue concentrations using BAF	Estimates amount of chemical that is taken up by resident species in the absence of site-specific data	BAF values are empirically derived and may not reflect actual conditions at the site
	Conduct bioaccumulation studies	Demonstrates whether metals in site soils/sediments are available for biological uptake	Bioaccumulation tests may be costly and more time consuming
Uptake from Food			
When evaluating absorption of metals from contaminated food	Perform laboratory bioassay to determine relative bioavailability	Provides measure of actual absorption of site-specific dose	Bioassays may be costly and time consuming

Direct Exposures to the Available Fraction. Estimates of the available fraction in sediment or soil can be determined analytically, using a variety of sequential extraction techniques (Tessier and Campbell, 1987; Campbell et al., 1988). Although no single extraction method can completely quantify the available fraction, use of a 1-N HCl extraction technique provides the best estimate (Luoma, 1989). Use of the metal concentration derived from this analytical technique as the EPC provides a more accurate estimate of the actual exposures to ecological receptors than the total metal concentration. As discussed in Section 2.3.2, in sediment these concentrations can be further refined to reflect consideration of AVS, which are operationally defined as the sulfide liberated from wet sediment by treatment with 1-N HCl (Ankley et al., 1996). Methods for applying this modification are described in Section 2.3.2.

In addition to the analytical determination of the bioavailable fraction, it is possible to qualitatively determine the potential for bioavailability based on certain chemical and physical parameters (e.g., pH, fraction organic carbon [f_{oc}], TOC, Eh). For example, adsorption of inorganic cations (e.g., Pb^{2+}) to soil increases with pH, with a resulting decrease in bioavailability, while the reverse is true for inorganic anions (e.g., $H_2AsO_4^{1-}$). Similarly, metals in sediments tend to be more bioavailable in acidic freshwater bodies than in neutral or basic waters. Seawater is naturally buffered at a pH of about 8.0 (alkaline), so most metals in marine sediments are less bioavailable than those in most freshwater systems. Based on this information, evaluation of soil pH can provide a quick, qualitative indication of whether measured metals are likely to be bioavailable. In addition, bioavailability and toxicity may vary depending on the form of the metal (see Section 6.0 of this document and U.S. EPA, 1992). Therefore, an understanding of the specific forms of the metal present also can assist in determinations regarding their potential bioavailability.

Toxicity tests of environmental media such as sediment and soil also can be used to evaluate the potential for bioavailability from environmental media. Typically, these tests are used to confirm assumptions made based on qualitative evaluations of chemical and physical parameters at the site. Although such tests do not provide a numerical estimate of the bioavailable fraction, the presence or absence of toxicity in organisms exposed to site materials versus reference materials provides an additional line-of-evidence argument for or against bioavailability. The combination of qualitative evidence indicating limited bioavailability and bioassays exhibiting low toxicity has been used successfully to demonstrate that metals at a site are not bioavailable.

Bioaccumulation from Environmental Media. Uptake and retention of metals by organisms (i.e., bioaccumulation) either may be measured directly by collecting and analyzing the tissues of representative organisms, or it may be estimated (BJC, 1998). As previously discussed, estimates of tissue concentration are derived by multiplying the concentration in environmental media (i.e., soil, sediment, or water) by a chemical-specific BAF typically found in the literature. Alternatively, BAF values can be derived from tissue and soil or sediment data available in the literature or even determined experimentally at the site. Determination of site-specific BAF values requires correlated concentrations in sediment or soil and tissues to provide an accurate representation.

Bioaccumulation of metals also may be evaluated through the use of bioaccumulation assays. These studies involve exposure of relevant species not previously exposed to metals to sediments or soils collected from the site. At the end of the test, the concentrations of metals in the tissues of the organism are determined. For the purpose of the bioassay, lower accumulation of metals from site soils or sediments relative to a reference material would indicate limited bioavailability at the site. Similar to toxicity studies, these bioassays may be used in the latter stages of an ecological risk assessment to provide an additional line of evidence regarding assumptions based on more qualitative approaches earlier in the process.

Uptake from Food. As discussed in Section 2.3.2, estimates of the uptake of metals from food by ecological receptors may be made using an ABS. However, identifying the appropriate ABS for use in an ecological assessment can be a complicated process. Section 2.1.1 describes the concept of relative bioavailability, which is used to derive RAFs for human health assessments. Although not typically considered for ecological assessments, this approach could be applied in the same manner to estimate the fraction of metal in food available to ecological receptors. To apply this approach to ecological assessments, it is suggested that the tests be designed to incorporate species representative of the key receptors identified at the site.

4.2 Steps in Conducting a Bioavailability Study

The key steps in conducting a bioavailability study are outlined in Figure 4-4. These steps apply mainly to human health bioavailability studies; however, they also can be used to guide bioavailability studies for ecological risk assessments, particularly if animal feeding studies are involved. As discussed in Section 3.1, bioavailability studies are typically done during the second tier of the risk assessment process. There are several factors in the figure that should be emphasized. First, it is important to thoroughly evaluate whether a bioavailability study is appropriate and feasible for the site before the study is undertaken (see Section 3.3). The key question that must be answered is whether a bioavailability study is likely to result in an adjustment to the risk estimate or cleanup goals that either will reduce the cost of remediation sufficiently to justify the increased cost and time required for the study, or will resolve another issue of concern such as avoiding impacts from remediation. Second, in development of the work plan, it is important to consider factors that will support the credibility of the study results, such as involving a qualified peer reviewer in development of the work plan, collecting representative samples, using accepted GLPs or the equivalent, and selecting a reputable testing laboratory. Finally, one of the most important factors is involving the regulators, and possibly other stakeholders, at the outset and giving them the opportunity to provide input throughout the process. By involving them early and giving them the opportunity for input along the way, they are more likely to accept the results. On the other hand, if they are not receptive to the concept of bioavailability adjustments, it is best to find this out early, before time and money are spent on bioavailability studies.

4.3 General Considerations

This section provides general information on cost, timing, and resources required to perform various test methods associated with a bioavailability study. It should be noted that the cost to run a study is only part of the total cost of successfully incorporating such studies into a site investigation. Cost and time for other components of the bioavailability study (e.g., developing a work plan and testing protocols, peer review of protocols and study results, and negotiations with regulators) need to be considered in planning the project.

4.3.1 Human Health Risk Assessments

Tables 4-2 and 4-3 present a summary of technical resources for conducting both in vitro and in vivo studies to estimate the relative bioavailability of metals from soil. Because in vitro methods are relatively well established for arsenic and lead, it is appropriate to perform these studies in commercial laboratories, and contact information is provided for several laboratories that have performed these types of tests. However, any competent analytical laboratory should be capable of performing these tests, and the listing of any particular laboratory in Tables 4-2 and 4-3 does not constitute an endorsement of that laboratory.




Figure 4-4. Steps in Conducting a Bioavailability Study

Table 4-2. Technical Resources for Conducting Bioavailability Studies for Use in Human Health Risk Assessments

Studies	Animal Model	Time Required	Cost	Contracting Institution^(a)
In Vitro (oral)				
Arsenic, cadmium, chromium, lead, and nickel (data only)	NA	3 weeks ^(b)	\$150/sample ^(c)	Univ. of Colorado at Boulder, CO ACZ Laboratories, Inc., Steamboat Springs, CO Bureau of Reclamation, Denver, CO
Arsenic, lead, cadmium, chromium, nickel, and mercury (full study) ^(d)	NA	6-8 weeks	\$5,000-15,000/study ^(e)	Qualified consulting firms
In Vivo (oral)				
Arsenic	Monkeys	3-6 months ^(f)	\$50-80,000/substrate ^(g)	Battelle Memorial Institute Univ. of Florida
Lead	Rats	3-6 months ^(f)	\$60-85,000/substrate ^(g)	Battelle Memorial Institute Univ. of Missouri
	Swine	3-6 months ^(f)	\$45,000/substrate ^(g)	
Cadmium	Rats	3-6 months ^(f)	\$60-85,000/substrate ^(g)	Battelle Memorial Institute
Mercury	TBD	5-8 months ^(h)	\$75-100,000/substrate ⁽ⁱ⁾	–
Chromium	TBD	5-8 months ^(h)	\$60-85,000/substrate ⁽ⁱ⁾	–
Nickel	TBD	5-8 months ^(h)	\$60-85,000/substrate ⁽ⁱ⁾	–
Dermal Absorption				
Arsenic, cadmium, chromium, lead, mercury, and zinc	Monkeys	3 months ^(f)	\$45-55,000/substrate	Univ. of California at San Francisco

(a) The laboratories listed in this table are provided only as a source of information. This list does not constitute a recommendation or an endorsement of these organizations. Contact information is provided in Table 4-3.

(b) Assumes sample extraction, and two-week analytical turnaround on analysis of a single metal in the extract and the test soil.

(c) Average per sample cost for data production only at a commercial analytical laboratory.

(d) Includes protocol development, sample handling and testing, report, production, and limited negotiations with a regulatory agency (phone calls only).

(e) Actual cost depends on number of samples, and project specific requirements.

(f) Includes external review of existing protocol, study, and reporting, but no agency negotiations.

(g) Actual cost depends on laboratory that is conducting the study and study design.

(h) Includes protocol development and external review, study, and reporting. No agency negotiations included.

(i) Represents an approximate cost estimate. No such study has been conducted to date.

TBD = To be determined.

NA = Not applicable.

Table 4-3. Contact Information for Laboratories Performing Human Health Bioavailability Studies^{(a) (b)}

Name	Specialty	Address	Contact Information
Dr. Peter Grevatt ^(c)	Bioavailability study issues/data use	Office of Solid Waste and Emergency Response U.S. EPA (MC 5103) 401 M Street, SW Washington, DC 20460	Phone: (202) 260-3100 Fax: (202) 401-1496 Grevatt.peter@epa.gov
Dr. John Drexler	Extraction tests for all metals	Univ. of Colorado at Boulder Dept. of Geological Sciences Campus Box 250 Boulder, CO 80309	Phone: (303) 492-5251 Fax: (303) 735-4953 drexler@spot.colorado.edu
Mr. Jammie Sabin	Extraction tests for As and Pb	ACZ Laboratories, Inc. 30400 Downhill Dr. Steamboat Springs, CO 80487	Phone: (970) 879-6590 Fax: (970) 879-2216 jammies@acz.com
Mr. Kevin Kelly or Ms. Barbara Frost	Extraction tests for As and Pb	Environmental Chemistry Research Laboratory Bureau of Reclamation PO Box 25007, Bldg. 67 Denver Federal Center Denver, CO 80225	Phone: (303) 445-6327 kkelly@de.usbr.gov Phone: (303) 445-6327 bfrost@de.usbr.gov
Dr. Jerry Johnson	Oral in vivo tests	Battelle Memorial Institute 505 King Avenue Columbus, OH 43201	Phone: (614) 424-4499 Fax: (614) 424-3171 johnsojd@battelle.org
Dr. Steven Roberts	Oral in vivo test for As	Univ. of Florida Center for Environmental & Human Toxicology, Bldg. 472, Mowry Rd Box 110885, Gainesville, FL 32611	Phone: (352) 392-4700 x 5505 Fax: (352) 392-4707 sroberts.vetmed1@mail.health.ufl.edu
Dr. Stan Casteel	Oral in vivo test for As and Pb	Univ. of Missouri College of Veterinary Medicine 1600 East Rollins Columbia, MO 65205	Phone: (573) 882-6811 Fax: (573) 882-1411 casteels@missouri.edu
Dr. Ronald Wester	Dermal in vivo and in vitro tests for all metals	Univ. of California at San Francisco Dermatology Dept. PO Box 0989 San Francisco, CA 94143	Phone: (415) 476-2468 Fax: (415) 753-5304 rcwgx@itsa.ucsf.edu

- (a) The laboratories listed in this table are provided only as a source of information. This list does not constitute a recommendation or an endorsement of these organizations.
- (b) All of these contacts are primarily experimental specialists, not risk assessors. It is recommended that these persons be contacted to perform specific tests (in vivo or in vitro), but not for design of entire studies. This list is subject to change.
- (c) Dr. Grevatt is listed as a contact for general information from U.S. EPA, not as a testing laboratory.

In vitro studies for cadmium, chromium, and nickel are no more complicated than those for arsenic and lead, and the same laboratory references are therefore applicable. Mercury, on the other hand, is more complicated to work with due to the potential for elemental mercury to volatilize, and it is recommended that a consulting firm that has qualified specialists in mammalian toxicology, soil chemistry, and aqueous geochemistry be contacted to perform these types of studies.

For the in vitro evaluation of all these elements, the cost of conducting the extraction and analyzing the extract is only a fraction of total study cost, if the study includes protocol development, external review,

reporting, and negotiations with the appropriate regulatory agency. Although individual samples may cost only a few hundred dollars to process through the simplified lead protocol, at least five samples per site are typically evaluated, and whenever any more complicated protocols are developed the total cost of developing protocols, running the study, and preparing a report will likely cost \$5,000 to \$15,000. At the upper end, these studies also would include mineralogical analyses to support interpretation of the in vitro extraction test results. Typically, in vitro studies can be planned, run, and reported in 6 to 8 weeks.

As described in Section 5.0, in vivo studies have been conducted to determine the relative bioavailability of arsenic, cadmium, and lead in soil, and contact information for the laboratories that have performed these studies is provided in Tables 4-2 and 4-3. The costs for in vivo studies, including protocol development and report preparation, will range from \$50,000 to \$200,000 depending on study design and number of samples tested. A minimum of 3 months is needed to order animals, allow for a quarantine period once the animals are ordered, run the study, get samples analyzed (with quality assurance review), and prepare a preliminary report. In planning a site investigation, it would be more realistic to allow for a total of 6 months from protocol development and review to final study report.

Most contract toxicology laboratories should be capable of performing these types of studies, and contact information is provided for some qualified laboratories (this list is by no means exhaustive). Contract laboratories are also likely to routinely conduct studies in accordance with GLPs (see Section 4.1.2), but generally will be unfamiliar with handling soil samples. University laboratories may provide a lower cost alternative for conducting these studies, but generally do not follow GLPs to conduct studies. Because successful relative bioavailability studies have not been conducted for chromium, mercury, and nickel, the initiation of such a study will require development of a detailed study protocol, external peer-review of the protocol, and possibly one or more pilot studies to ensure that an appropriate animal model has been selected.

Because no dermal absorption studies have been conducted for soils that contain the forms of metals commonly found in the environment, undertaking such a study will require careful planning and execution. Dr. Ronald Wester, who is a research dermatologist at the University of California at San Francisco, performed the existing studies on the dermal absorption of soluble forms of arsenic, cadmium, and mercury in the presence of soil. Therefore, Dr. Wester's laboratory would be one that is qualified to perform the required studies on environmental soil samples (see Table 4-3 for contact information).

4.3.2 Ecological Risk Assessments

Table 4-4 provides a summary of the estimated cost and time for each of the different tests and analyses proposed for measuring bioavailability in ecological risk assessments. These costs are intended to provide an indication of the analytical level of effort necessary to address these issues and may not reflect actual total costs associated with each task. In general, all of the tests proposed are standard laboratory protocols for which specific methods have been developed. For example, ASTM publishes guidance on the appropriate methodologies for evaluating the toxicity of metals to aquatic and terrestrial invertebrates. Similarly, the analytical methods discussed rely on standard analytical techniques. As a result, these tests can be performed by any qualified laboratory. The cost estimates provided are averages for contract laboratories; other laboratory facilities (e.g., universities) may offer lower costs for some of these analyses.

It is important to note that the exact cost of a bioavailability study will vary from site to site, depending on the existing data and the complexity of the site. For example, if all chemical and physical parameters are available from existing data, it may not be necessary to collect additional samples. In addition, costs could not be estimated for qualitative evaluations (e.g., incorporation of a literature-based BAF), or for

interpretation of results or negotiations with agencies. It is impossible to accurately predict the costs associated with these tasks because their scope is entirely dependent on site-specific factors including the size of the site, tests selected for inclusion, and the technical expertise available to the Navy. In some instances, the Navy may require additional technical expertise for assistance in data interpretation, while at other sites, such assistance may not be required. Therefore, the costs in Table 4-4 are offered to provide a general background on the relative costs of the various tests proposed.

Table 4-4. Time and Cost Associated with Test Methods for Assessing Bioavailability in Ecological Risk Assessments

Test Type	Description	Estimated Cost per Sample^(a)	Time per Test
Direct Exposures to the Available Fraction			
Extraction Techniques	1-N HCl	\$120	Allow 3-4 weeks for sample analysis
Comparison of AVS/SEM	Compare ratio of measured SEM to AVS	\$250	Allow 3-4 weeks for sample analysis
Evaluation of chemical and physical parameters	Chemical form, pH, TOC, Eh, f_{oc} , etc.	\$200	Allow 3-4 weeks for sample analysis
Toxicity tests	Standard test methods for aquatic or terrestrial invertebrates	\$500-1,200	Test lengths can vary from 10 to 28 days
Bioaccumulation from Environmental Media			
Collect and analyze site-specific tissue data	Metals in fish, invertebrates, birds, mammals, etc.	\$300-400 ^(b)	Allow 3-4 weeks for sample analysis
Estimate using BAF	Literature-based (reported or derived); site-specific	Level of effort will vary depending on number of chemicals and species evaluated	Level of effort will vary depending on number of chemicals and species evaluated
Conduct bioaccumulation studies	Standard test methods for aquatic or terrestrial invertebrates	\$1,900 per species (includes cost of 5 replicates and chemical analyses)	Test lengths can vary from 10 to 28 days
Uptake from Food			
Relative bioavailability study	As described in Section 4.1.2 (also see Section 5.7)	In vivo: \$45,000-\$100,000/substrate (see Section 4.3.1)	In vivo: 3-6 months (see Section 4.3.1)

(a) Costs provided are estimated based on standard procedures. Total may vary depending on such factors as the specifics of project protocol and the number of chemicals analyzed.

(b) Costs provided assume analysis of whole body concentrations.

5.0 CHEMICAL-SPECIFIC CONSIDERATIONS FOR ASSESSING BIOAVAILABILITY TO HUMAN RECEPTORS IN TERRESTRIAL (SOIL) SETTINGS

This section provides a review of chemical-specific issues to consider when attempting to determine whether to proceed with site-specific bioavailability studies. The six metals included are those that are commonly important in human health risk assessments, specifically arsenic, cadmium, chromium, lead, mercury, and nickel. For each metal, the predominant forms in soil are briefly described. Differences in toxic endpoints for the different forms of the same metal are noted because evaluation of relative bioavailability is relevant only for forms of a metal that have the same toxic endpoints. The focus of the toxicity discussion is on oral toxicity. Generally, little or no toxicity data are available related to systemic effects of dermally applied metals. As described in section 4.1.2, it is unlikely that site-specific bioavailability studies of inhaled metals from resuspended soil particles will be useful. Consequently, inhalation toxicity and bioavailability of metals is excluded from this discussion.

For each metal, available data documenting variations in oral bioavailability from different media are described. Oral absorption of arsenic and lead from soil has been studied quite extensively and studies of cadmium and mercury, although limited, have been conducted. The oral bioavailability of chromium and nickel in soil is not well characterized. The database for dermal bioavailability is much more limited. Dermal absorption studies have been conducted for arsenic, cadmium, and mercury in soil, but in all three cases, soluble forms of the metals were mixed with soils and tested without time for weathering reactions to occur. Thus, there are no data currently available to predict the bioavailability of these metals in weathered soils at contaminated sites.

5.1 Arsenic

Default risk-based soil cleanup levels for arsenic are frequently below local background soil concentrations of this element. If cleanup levels in soil are based on background concentrations, site-specific bioavailability data may have a limited impact on cleanup levels when the adjusted risk-based cleanup levels are still below background concentrations. Nevertheless, in situations where there is some flexibility in target risks, bioavailability data may be a powerful tool for adjusting cleanup goals.

5.1.1 Predominant Forms in Soil

Trivalent and pentavalent inorganic arsenic compounds are the predominant forms in soils. Inorganic arsenic compounds vary widely in their water solubility, with sodium arsenate and arsenic trioxide representing highly water-soluble forms. Discrete arsenic mineral phases present in soils commonly include less soluble forms such as sulfide minerals, complex oxides, and arsenic present in iron, manganese, and phosphate mineral species. All but the sulfide minerals may be formed over time in surficial (oxygen-rich) soils, as weathering reactions occur that favor the most thermodynamically stable metal forms. Arsenic may also be present in soil in ionic forms that may be adsorbed to soil constituents. Reduced bioavailability of arsenic in soil is thought to be primarily a function of the presence of less soluble mineral phases and ionic forms that are strongly adsorbed to soil particles or coprecipitated with other elements in soil.

5.1.2 Toxicity Assessment

All inorganic arsenic compounds induce chronic toxic effects by the same mechanism, regardless of valence state. Ingested inorganic arsenic compounds cause cancer at high doses, so all inorganic arsenic compounds may be considered together when assessing bioavailability. The oral toxicity values used in

risk assessments are based on epidemiology studies of human populations exposed to soluble inorganic arsenic dissolved in drinking water, so these soluble forms should be the point of comparison in studies of relative bioavailability.

5.1.3 Relative Bioavailability Via Oral Exposure

After ingestion, water-soluble forms of inorganic arsenic are almost completely absorbed from the gastrointestinal tract of humans and many laboratory animals. Ingestion of less soluble forms of arsenic leads to reduced absorption. Studies have been conducted in laboratory animals that demonstrate reduced absorption of arsenic from soil taken from many different sites (Freeman et al., 1993; Freeman et al., 1995; Groen et al., 1994; Casteel et al., 1997b; Rodriguez et al., 1999). These studies indicate that arsenic in soil is typically only one-half to one-tenth as bioavailable as soluble arsenic forms. In other words, these studies support relative bioavailability adjustments ranging from 0.5 to 0.1 in exposure assessments for these sites.

Monkeys, dogs, rabbits, and swine have been used to study arsenic in soil, mainly from mining and smelting sites. Bioavailability estimates have been based on the fraction of the dose excreted in the urine, and on the AUC values for arsenic concentrations in the blood. Figure 5-1 illustrates differences in excretion of soluble arsenic and arsenic from soil and indoor dust from Anaconda, MT in the urine of monkeys. The animal studies are supported by mineralogical analyses demonstrating the presence of less soluble arsenic forms in the soils tested, and by in vitro studies (i.e., PBETs) that indicate reduced bioaccessibility of arsenic in the samples studied.




Figure 5-1. Monkey Bioavailability Study: Arsenic Excretion in Urine

5.1.4 Bioavailability Via Dermal Exposure

The dermal bioavailability of a water-soluble arsenic form (sodium arsenate) mixed with a soil matrix has been evaluated in vivo in monkeys, yielding estimates of arsenic absorption from soil ranging from 3.2 to 4.5 percent (Wester et al., 1993a). The same soil mixture was tested with human skin in vitro, yielding an estimate of approximately 1 percent absorption. As a result of this study, a value of 3 percent dermal absorption of arsenic from soil is being used in some risk assessments.

5.1.5 Summary of Pertinent Data

Inorganic forms of arsenic vary in water solubility and bioavailability. Most of the oral bioavailability studies of soil arsenic conducted to date used soil from mining or smelting sites, and support relative bioavailability adjustments ranging from 0.5 to 0.1. A simple in vitro test system is available that has shown good agreement with the results of studies in laboratory animals using the same soils (Rodriguez et al., 1999; see Section 4.1.1).

5.2 Cadmium

Risk-based soil cleanup levels for cadmium may be influenced by dermal exposures and by uptake into homegrown produce, as well as by direct ingestion of soil. Therefore, the relative importance of these pathways should be evaluated prior to planning site-specific bioavailability studies.

5.2.1 Predominant Forms in Soil

Cadmium in soil may be found in forms that range in solubility from sparingly (sulfides) to moderately (cadmium sulfate) to highly soluble (cadmium carbonate).

5.2.2 Toxicity Assessment

The reference dose for cadmium is based on effects of a soluble form of cadmium (cadmium chloride) on the kidney. All inorganic cadmium forms commonly present in soils induce chronic toxic effects after ingestion by the same mechanism. Consequently, all inorganic cadmium compounds may be considered together when assessing bioavailability.

5.2.3 Relative Bioavailability Via Oral Exposure

Oral absorption of cadmium in humans generally is reported to be very low (1 to 7 percent) (ATSDR, 1997a). Evidence that the bioavailability of cadmium in soil may be reduced compared to the bioavailability of soluble cadmium forms is available from a limited number of studies. Several studies have reported reduced oral bioavailability of a soluble cadmium form, cadmium chloride, mixed with soil (Griffin et al., 1990; Schilderman et al., 1997). For cadmium in weathered soil, data are available for soil from a single site (the site of a former zinc smelter) that has been evaluated in vivo in rats (Schoof and Freeman, 1995; PTI, 1994). A relative cadmium bioavailability estimate of 33 percent was obtained based on comparison of liver and kidney tissue concentrations in animals fed rodent chow mixed with soil, versus those fed rodent chow mixed with cadmium chloride. An in vitro study of this same soil yielded a higher value, which suggests that the in vitro method might overestimate the relative bioavailability of soil cadmium.

5.2.4 Bioavailability Via Dermal Exposure

An in vitro study of dermal absorption in human cadaver skin of cadmium chloride mixed with soil yielded an estimate of 0.02 to 0.07 percent absorption based on cadmium in receptor fluid (Wester et al., 1992). An additional 0.06 to 0.13 percent of the dose was retained in the skin. The U.S. EPA default value of 1.0 percent for dermal absorption of cadmium compounds from soil is more than 10 times higher than the maximum percent of the cadmium chloride dose reaching the receptor fluid and 5 times higher than the maximum combined percent dose in receptor fluid and skin. Dermal absorption of cadmium from weathered soils may be even lower.

5.2.5 Summary of Pertinent Data

Limited evidence is available that oral absorption of cadmium in soil is reduced compared to absorption of soluble cadmium. For any site in which dermal exposures are quantified, the highest priority for site-specific studies may be studies of dermal exposure from soil. This priority reflects the likelihood that default assumptions overestimate dermal absorption of cadmium from soil by a factor of 10 or more, but may only overestimate oral absorption by a factor of 3.

5.3 Chromium

The two primary oxidation states of chromium are trivalent and hexavalent, with hexavalent chromium generally being more bioavailable and more toxic than trivalent chromium. Sometimes soil cleanup levels for total chromium are based on the toxicity value for hexavalent chromium. In such cases, it clearly would be prudent to characterize the form of chromium present before trying to decide if bioavailability studies would be useful.

5.3.1 Predominant Forms in Soil

Unlike many of the other metals discussed in this document (e.g., arsenic, cadmium, and lead), anthropogenic sources of chromium for soils are generally in a soluble form (with the exception of sites that contain chromite ore processing residue). As a result, the soil alteration processes that control chromium bioavailability generally have these soluble chromium species as a starting point. The solubility and mobility of trivalent chromium is minimal, whereas hexavalent chromium is both highly soluble and mobile. The relative concentrations of trivalent chromium and hexavalent chromium in a particular soil sample will depend on the form of the chromium contaminant and the soil redox conditions and geochemistry, particularly the pH and presence of oxidizing or reducing agents.

5.3.2 Toxicity Assessment

Trivalent chromium is a required nutrient. The oral reference dose for trivalent chromium applies to insoluble salts, and is based on a study in which no adverse effects were observed at any dose tested when Cr_2O_3 was baked into bread and fed to rats. The oral reference dose for hexavalent chromium applies to soluble salts, and is based on doses that caused no adverse effects in a rat drinking water study. Based on their respective reference doses, soluble salts of hexavalent chromium are considered to be almost 1,000 times more toxic than insoluble salts of trivalent chromium.

5.3.3 Relative Bioavailability Via Oral Exposure

The oral bioavailability of chromium depends on its valence state, with hexavalent chromium being more readily absorbed than trivalent chromium (ATSDR, 1998). Oral absorption of nondietary trivalent

chromium compounds is extremely low (approximately 1 percent). Absorption of hexavalent chromium compounds is somewhat higher (approximately 10 percent). There is evidence that hexavalent chromium is converted to trivalent chromium in the acid environment of the stomach, which would limit the oral bioavailability of hexavalent chromium. Two oral in vivo studies using environmental soil chromium samples are reported in the literature, one performed in humans and one in laboratory animals (Gargas et al., 1994; Witmer et al., 1989, 1991). Both studies used soils containing chromite ore processing residues, and therefore contained a mixture of trivalent and hexavalent chromium. Although these studies suggested limited oral absorption of the soil chromium, no reliable estimates of relative bioavailability were obtained.

5.3.4 Bioavailability Via Dermal Exposure

Hexavalent chromium and trivalent chromium exhibit very limited ability to penetrate the skin, with somewhat greater penetration observed for hexavalent chromium. Less than 1 percent absorption of hexavalent chromium from water was observed for dosing periods of 5 hours (Wahlberg and Skog, 1963). No studies of dermal absorption of chromium from soil were identified.

5.3.5 Summary of Pertinent Data

The complexity of the factors affecting chromium geochemistry combined with differences in toxicity make it necessary to characterize the valence states of chromium in soils at a site prior to beginning any site-specific bioavailability studies.

5.4 Lead

Direct ingestion of lead in soil and dust generally drives soil lead cleanup levels. Lead is the only chemical for which the U.S. EPA's default assumption is that oral bioavailability from soil is less than the oral bioavailability of soluble forms. Methods for assessing the oral bioavailability of lead in soil are well developed, and are relatively easy to conduct on a site-specific basis.

5.4.1 Predominant Forms in Soil

Inorganic lead is present in geologic materials and soils in more than 200 minerals that vary greatly in solubility. The majority of lead in geologic materials is in the form of galena (lead sulfide), anglesite (lead sulfate), and cerussite (lead carbonate). Organic forms of lead are rare in soils and are not evaluated in this document.

5.4.2 Toxicity Assessment

The toxicity assessment for lead used by the U.S. EPA is unique, incorporating specific assumptions for lead absorption from ingested water, food, and soil in a pharmacokinetic model that predicts lead levels in blood. Inorganic forms of lead in soil all have the same toxic endpoints and may be considered together when assessing bioavailability.

5.4.3 Relative Bioavailability Via Oral Exposure

Gastrointestinal absorption of lead varies with the age, diet, and nutritional status of the subject, as well as with the chemical species and the particle size of lead that is administered (ATSDR, 1993b). Age is a well-established determinant of lead absorption; adults typically absorb 7 to 15 percent of lead ingested from dietary sources, while estimates of lead absorption from dietary sources in infants and children range

from 40 to 53 percent. In U.S. EPA's childhood lead model, it is assumed that 50 percent of an oral lead dose is absorbed from food and water, while 30 percent of a soil lead dose is assumed to be absorbed. Thus, the default assumption for lead is that the relative bioavailability of soil lead compared to soluble lead forms is 0.6 (i.e., 30 percent divided by 50 percent) (U.S. EPA, 1994a).

The oral bioavailability of lead in soil has been more extensively studied than that of any other metal. Soil lead absorption has been studied in rats, swine, and humans. The swine model has been used to test soils from numerous sites. A physiologically based extraction method is also well developed (Ruby et al., 1993, 1996; Medlin, 1997) and is undergoing detailed validation studies.

The studies in rats and swine have indicated that absorption of lead from soil will vary with the source of the lead, ranging from near zero to greater than 50 percent absolute bioavailability (i.e., relative bioavailability of 1.0, or more compared to soluble lead forms) (Casteel et al., 1997a; Dieter et al., 1993; Freeman et al., 1992, 1996a; Schoof et al., 1995; U.S. EPA, 1996b-e; 1998a-e). On average, the results of these studies support the use of a default assumption that 30 percent of an oral lead dose is absorbed from soil (i.e., relative bioavailability of 0.6). A study in adult humans indicates that absolute lead bioavailability from a mining-area soil varies from approximately 3 to 26 percent, depending on how recently the test subject had eaten (Maddaloni et al., 1998).

5.4.4 Bioavailability Via Dermal Exposure

It is generally assumed that absorption of inorganic lead compounds through the skin is negligible in comparison to the oral or inhalation routes, and dermal exposure to soil lead is generally excluded from risk assessments. No studies of the dermal absorption of lead from soil or dust were identified.

5.4.5 Summary of Pertinent Data

A substantial body of research has demonstrated that the relative oral bioavailability of soil lead varies from site to site. On average, the current default assumption that the relative oral bioavailability of soil lead is 0.6 has been found to be appropriate. A simple in vitro extraction method, currently being validated for lead, may offer a rapid, cost-effective method for generating site-specific data.

5.5 Mercury

Mercury is the only metal for which inhalation of vapors released from soil may be an exposure pathway of concern. If elemental mercury is present in soils at a site, the relative importance of the inhalation exposures compared to oral exposures should be assessed prior to determining whether oral or dermal bioavailability studies would be useful.

5.5.1 Predominant Forms in Soil

Mercury in contaminated soils generally is present as either elemental mercury or inorganic mercury compounds. Organic mercury compounds are rarely present in soil in significant quantities. Consequently, only the inorganic forms of mercury are considered here. Inorganic mercury species in weathered soils range from forms with extremely limited solubility (i.e., elemental mercury and mercuric sulfide) to much more soluble forms (i.e., mercury adsorbed into organic matter or clays, and mercury oxides, hydroxides, and carbonates).

5.5.2 Toxicity Assessment

Because of significant differences in pharmacokinetic characteristics and toxicity, elemental mercury and all other inorganic mercury compounds must be addressed separately. The oral reference dose typically applied to inorganic mercury compounds is specifically described as a reference dose for mercuric chloride, a water soluble form of mercury. This reference dose is based on autoimmune effects observed in rats. There is no oral reference dose for elemental mercury due to its extremely limited oral absorption. However, if elemental mercury is present in surface soils, risk-based cleanup levels will be driven by predicted inhalation exposures from mercury vapor released from soil.

5.5.3 Relative Bioavailability Via Oral Exposures

Soluble forms of inorganic mercury, such as mercuric chloride or mercuric nitrate, appear to be 15 to 25 percent absorbed across the gastrointestinal tract (Rahola et al., 1973; Nielsen and Anderson, 1990). Several studies suggest that mercuric sulfide, a relatively insoluble inorganic mercury compound, has a much lower bioavailability than mercuric chloride (i.e., approximately 30 to 60 times lower) (Schoof and Nielsen, 1997). The oral absorption of elemental mercury is quite low, perhaps on the order of 0.01 to 0.1 percent (ATSDR, 1997b).

One study has been identified that attempted to estimate the bioavailability of mercury in environmental soil samples using an animal model (Revis et al., 1989, 1990), but the study did not yield reliable bioavailability estimates because of study design limitations. Another study suggests that the presence of soil alone decreases the oral bioavailability of inorganic mercury compounds (Sheppard et al., 1995). Several in vitro studies performed to measure the dissolution of mercury from soil found that relative bioavailability was generally estimated to be less than 10 percent (SAIC, 1994; CDM, 1992).

5.5.4 Bioavailability Via Dermal Exposure

A study of dermal absorption of mercuric chloride from water and soil used an in vitro model with human cadaver skin (Wester et al., 1995). In this study, very little mercury passed through the skin and appeared in the receptor fluid (0.7 percent for water, 0.06 percent for soil), but a substantial amount of mercury was retained in the skin (28.5 percent for water, 7.9 percent for soil). It is not clear what proportion of the mercury retained in the skin would subsequently be absorbed.

5.5.5 Summary of Pertinent Data

Due to differences in toxicity and predominant routes of exposure, it is necessary to identify the mercury species present in soil whenever bioavailability studies are performed. Speciation studies of mercury are technically challenging, and peer review of proposed methods is recommended. Studies of oral absorption of mercury from weathered soils are very limited, and no dermal absorption studies have used weathered soils.

5.6 Nickel

Little is known about the bioavailability of nickel compounds in soil. Due to the low oral absorption of nickel compounds, predicted dermal exposures may be significant if included in risk assessments. Consequently, both oral and dermal bioavailability studies may be needed.

5.6.1 Predominant Forms in Soil

Nickel may be present in soils in a variety of mineral forms, from forms with very limited solubility (sulfide and sulfate forms) to the much more soluble carbonate form. Given that nickel may be present as discrete mineral phases of varying solubility in soils, or adsorbed onto organic matter or clay particles, the solubility of nickel in soils will vary with different nickel sources and soil geochemistry.

5.6.2 Toxicity Assessment

The nature of the oral toxicity of nickel does not vary among the different forms expected to be present in soil. The oral reference dose is based on a study in which a soluble nickel salt (nickel sulfate hexahydrate) administered to rats after being mixed with their diet caused reduced body and organ weights. Roughly 10 to 15 percent of the population will show an immunological contact dermatitis reaction in response to nickel applied to the skin (Peltonen, 1979). This localized effect will not be dependent on systemic absorption, but may be affected by the solubility of nickel forms contacting the skin.

5.6.3 Relative Bioavailability Via Oral Exposures

Nickel generally is not well absorbed from the gastrointestinal tract in either laboratory animals or humans (ATSDR, 1997c). Less than 5 percent of the most soluble nickel salts are absorbed orally in humans and animals. The gastrointestinal absorption of nickel correlates directly with the solubility of the metal, with less than 1 percent of the least soluble forms (oxides and sulfides) being absorbed.

When a soluble nickel form, nickel chloride, was mixed with soil and administered to rats as an aqueous slurry, the bioavailability was reduced relative to nickel chloride administered to the rats in water (Griffin et al., 1990). The sandy-loam slurry produced a relative bioavailability of 63.1 percent, and the clay-loam slurry a 33.5 percent relative bioavailability, as measured by nickel in blood. No studies of the relative oral bioavailability of nickel in weathered soils were identified.

5.6.4 Bioavailability Via Dermal Exposures

No studies of dermal absorption of nickel from soil were identified.

5.6.5 Summary of Pertinent Data

Because of the great variation in solubility of nickel compounds, site-specific studies of the relative oral bioavailability of nickel in soil could have a significant effect on risk-based cleanup levels.

5.7 Relevance to Ecological Receptors in Terrestrial Settings

All of the metal-specific considerations discussed above for assessing bioavailability to human receptors are applicable to certain terrestrial ecological receptors that are exposed to metals in soils through direct contact. However, direct comparisons are limited to monogastric mammalian receptors (e.g., small mammals and other wildlife), and do not necessarily apply to ruminants (e.g., deer or cows), reptiles, amphibians, and avian species. Small mammals that burrow in soils and exhibit preening behavior, or that ingest earthworms for a large portion of their diet, have elevated soil ingestion rates. For example, short-tailed shrew and eastern cottontail rabbits are estimated to consume 13 and 6.3 percent soil in their diet, respectively (Talmage and Walton, 1993; Sample and Suter, 1994). As a result, these receptors often drive ecological risk assessments for metals in upland soils. Because the TRVs used in ecological risk

assessment are based on laboratory studies where soluble metal salts were added to the diet of these animals, relative oral bioavailability becomes an important issue. When attempting to evaluate the importance of metals bioavailability from soil to these receptors, the metal-specific oral bioavailability values discussed above are applicable.

6.0 CHEMICAL-SPECIFIC CONSIDERATIONS FOR ASSESSING BIOAVAILABILITY TO ECOLOGICAL RECEPTORS IN AQUATIC (SEDIMENT) SETTINGS

All sediments contain metals. The metals in freshwater and marine sediments originate from several natural and human sources and are present in the sediments in several different physical and chemical forms (Goldberg, 1954). The chemical species and forms of complexed, adsorbed, and solid metals in sediments have a profound effect on the bioavailability and toxicity of the metals to aquatic/marine plants and animals (Nelson and Donkin, 1985). Each metal has unique physical and chemical properties that determine the forms of the metals in sediments and pore water and their relative bioavailability to aquatic receptors. Metals in highly insoluble solid forms are not bioavailable to sediment-dwelling organisms. Metals in solution or colloidal suspension in sediment pore water or in adsorbed forms that are readily desorbed (leached) into the dissolved phase by small changes in oxygen concentration, pH, and Eh are bioavailable. Therefore, it is important to understand the chemical forms of metals in sediments if bioavailability is going to be used in ecological risk assessment. The sections that follow are a brief summary of the forms, bioavailability, and toxicity of several metals in sediments.

Table 6-1 summarizes information on background concentrations and effects levels for the metals discussed in this section with the exception of tin. In addition, “high” concentrations developed by Daskalakis and O’Connor (1995) based on data from the National Oceanic and Atmospheric Administration’s (NOAA’s) National Status and Trends Program are included. Daskalakis and O’Connor (1995) examined chemical residue data for large numbers of marine sediment samples collected as part of the NOAA National Status and Trends Program and several other monitoring programs in coastal marine environments in the United States. They defined a “high” concentration of chemicals in sediments as the geometric mean concentration plus one standard deviation of the National Status and Trends site means.

**Table 6-1. Typical Background Concentrations and “High”
Concentrations of Metals in Coastal Sediments**

Metal	Background Conc. (µg/g dry wt)	High Conc. (µg/g)	ERL ^(a) (µg/g)	ERM ^(a) (µg/g)	Acute/Chronic Water Quality Criteria (µg/L)
Arsenic (As)	5-15	13	8.2	70	69/36
Cadmium (Cd)	0.1-0.6	0.54	1.2	9.6	43/9.3
Chromium (Cr)	50-100	125	81	370	1,100/50
Copper (Cu)	10-50	42	34	270	4.8/3.1
Lead (Pb)	5-30	45	46.7	218	220/8.5
Mercury (Hg)	≤ 0.2	0.22	0.15	0.71	2.1/1.11 ^(b)
Nickel (Ni)	≤ 50	42	20.9	51.6	75/8.3
Zinc (Zn)	1.2->100	135	150	410	95/86

(a) Effects Range Low (ERL) and Effects Range Median (ERM) Screening Levels for Marine Sediments and Acute/Chronic Marine Water Quality Criteria are Included

(b) Marine water quality values are for inorganic mercury. The chronic value of methylmercury is 0.025 µg/L.

6.1 Arsenic

6.1.1 Predominant Forms in Sediment

Concentrations of total arsenic in uncontaminated nearshore estuarine and marine sediments usually fall in the range of 5 to 15 $\mu\text{g/g}$ dry wt (Neff, 1997a) (Table 6-1). Daskalakis and O'Connor (1995) defined a "high" concentration of chemicals in sediments as the geometric mean concentration plus one standard deviation of the National Status and Trends site means. The "high" concentration of arsenic in coastal sediments is 13 $\mu\text{g/g}$. This concentration is exceeded frequently in sediments near natural (e.g., phosphate deposits) and anthropogenic sources of this chemical.

Arsenate (+V) is the most abundant form of arsenic in oxidized marine sediments, whereas arsenite (+III) is the dominant dissolved and solid species in reduced sediment layers (Neff, 1997a) (Table 2-3). Arsenite in oxidized sediments is oxidized rapidly to arsenate (De Vitre et al., 1991). Much of the arsenic in the oxidized layers of sediment is associated (coprecipitated or adsorbed) with the hydrous iron and manganese oxide fraction or is present as $\text{Fe}_3(\text{AsO}_4)$. Under these conditions, the amount of arsenic in solution in potentially bioavailable forms in oxidized sediment pore water is low and 65 to 98 percent is present as the less bioavailable arsenate (Masscheleyn et al., 1991).

Under moderately reducing conditions, iron and manganese oxide phases begin to dissolve, releasing adsorbed arsenate into pore water (Masscheleyn et al., 1991). Arsenate is reduced to arsenite in reducing sediments and, if sulfur is abundant (as is the case in most marine sediments), most of the arsenic reacts with sulfides to form realgar (AsS), impurities in copper and zinc sulfides, arsenopyrite (FeAsS), and orpiment (As_2S_3) (Morse, 1994). These sulfides have low solubility, mobility, and bioavailability.

However, in estuarine and freshwater sediments containing low concentrations of sulfur, arsenic solubility is less limited by formation of insoluble sulfide minerals. Arsenite, often as arsenolite (As_2O_3), may remain quite mobile and tends to diffuse upward to be released into the overlying water column as either arsenate or arsenite (Soma et al., 1994). Because of this behavior, the bioavailability of arsenic usually is highest in freshwater sediments, is intermediate in estuarine sediments, and is lowest in marine sediments.

6.1.2 Bioavailability and Toxicity in Sediments

Sediments are a major source of arsenic in bottom-living freshwater and marine animals (Bryan and Langston, 1992). There is a direct relationship between the concentration of arsenic in tissues of sediment invertebrates and the arsenic/iron (As/Fe) ratio in the easily extractable (1-N HCl) fraction of sediments in which the invertebrates reside. In uncontaminated or slightly contaminated oxidized sediments, most of the non-residual arsenic is adsorbed to iron oxyhydroxides and is relatively unavailable.

Concentrations of total arsenic in the tissues of marine invertebrates and fish are very high. Most of the arsenic is present as various organo-arsenic compounds, particularly arsenobetaine, which are not toxic to the marine animals or their consumers, including humans (Neff, 1997a).

Inorganic arsenic is more toxic to aquatic plants than aquatic animals. Arsenite and arsenate have similar toxicities to aquatic organisms, but different species differ markedly in sensitivity to arsenic (Neff, 1997a). Methyl-arsenic compounds, frequently present at trace concentrations in sediments, are bioavailable, but have a low toxicity. The U.S. EPA acute and chronic water quality criteria for arsenic (as arsenite) for protection of marine life are 69 $\mu\text{g/L}$ and 36 $\mu\text{g/L}$, respectively (Table 6-1). ERL and ERM concentrations of arsenic in marine sediments are 8.2 $\mu\text{g/g}$ and 70 $\mu\text{g/g}$, respectively (Long et al., 1995). Concentrations below the ERL values are considered to be rarely, if ever, toxic to bottom-

dwelling marine animals. Concentrations between the ERL and ERM may be toxic to some species. Concentrations above the ERM are nearly always toxic to most species.

6.2 Cadmium

6.2.1 Predominant Forms in Sediment

Cadmium concentrations in uncontaminated marine sediments usually are in the range of 0.1 to 0.6 µg/g dry wt (Warren, 1981) (Table 6-1). The “high” concentration of cadmium in coastal sediments is 0.54 µg/g (Daskalakis and O’Connor, 1995). There is a direct correlation in relatively uncontaminated sediments between concentrations of cadmium and aluminum (an indicator of clay minerals) (Schropp et al., 1990).

Cadmium in oxidized sediments is associated primarily (50 to 70 percent) with the carbonate plus iron/manganese oxide fractions of the sediment (Rosental et al., 1986) (Table 2-3). Most of the remainder is associated with the organic/sulfide fraction. Only about 1 percent is in the completely non-bioavailable residual fraction, indicating that cadmium associated with oxidized sediments is likely to be moderately mobile and bioavailable (Samant et al., 1990).

Cadmium in anoxic sediments appears to be associated almost exclusively with the sulfide phase (Salomons et al., 1987). Cadmium forms solid sulfides and strong complexes with sulfides. However, soluble cadmium sulfide complexes are formed (e.g., $\text{Cd}(\text{HS})_x^{x-2}$ where $x = 1$ or 4) only at high concentrations of sulfide ($>10^{-3}$ M). Cadmium sulfide complexes are moderately soluble; therefore, the mobility of cadmium in reducing environments may be quite high (Boulègue, 1983). Various insoluble hydroxide complexes may be present in freshwater sediments containing low sulfide concentrations. Nearly 90 percent of the cadmium in anoxic marine sediments is present as cadmium sulfide (Lee and Kittrick, 1984).

6.2.2 Bioavailability and Toxicity in Sediments

Marine invertebrates and fish bioaccumulate cadmium primarily from food and sediments (Canli and Furness, 1995; Wen-Xiong and Fisher, 1996). Oysters are able to filter 85 to 95 percent of cadmium-contaminated particles (sediment and diatoms) from water and retain about 60 percent of the cadmium supplied (Hardy et al., 1984). More than half the cadmium in the oyster tissues is from ingested particles; the rest is from bioconcentration from the water. When mice are fed cadmium-contaminated oysters, they retain about 0.83 percent of the administered dose in their tissues (Sullivan et al., 1984). Thus, the trophic transfer of cadmium from sediment particles and primary producers to a primary consumer is moderately efficient, but transfer to a secondary consumer, the mouse, is inefficient. Cadmium is not biomagnified in aquatic food webs.

Cadmium in ionic, bioavailable forms is one of the more toxic metals to freshwater and marine animals (Eisler, 1985). Toxicity tends to decrease with increasing salinity, because of complexation of the toxic ionic species with chloride. The U.S. EPA acute and chronic marine water quality criteria for cadmium are 43 µg/L and 9.3 µg/L, respectively (Table 6-1). ERM and ERL values for cadmium in sediments are 1.2 µg/g and 9.6 µg/g, respectively (Long et al., 1995).

6.3 Chromium

6.3.1 Predominant Forms in Sediment

Concentrations of total chromium in uncontaminated estuarine and marine sediments usually are in the range of 50 to 100 $\mu\text{g/g}$ dry wt (Mayer, 1988) (Table 6-1). The “high” concentration of chromium in U.S. coastal sediments is 125 $\mu\text{g/g}$ (Daskalakis and O’Connor, 1995). Much of the chromium in sediments is associated with the clay fraction, as indicated by a close correlation between aluminum and chromium concentrations (Schropp et al., 1990).

The distribution of chromium in sediment seems to depend in part on the source of the chromium. Generally, chromic chromium (+III) is more abundant than chromate chromium (+VI) in sediments. Chromate is a strong oxidizing agent and is reduced rapidly by organic matter and some metals in sediments. The small amounts of chromate in sediments usually is tightly bound to soil organic matter and iron oxide coatings on clay particles, or is coprecipitated with iron sulfides (Olazabal et al., 1997). In estuaries receiving chromium from tanneries and electroplating operations, more than 80 percent of the total chromium in the sediment is associated with the organic/sulfide fraction (Loutit et al., 1988). Because chromium is not known to form sulfides, carbonates, or phosphates (Mayer, 1988), and because of the stability of solid $\text{Cr}(\text{OH})_3$, it is probable that most of the chromium in these sediments is bound to organic matter or is present as the hydroxide (Table 2-3).

Chromium in less contaminated oxidized sediments often is adsorbed primarily to amorphous iron oxide (50 to 70 percent) and organic/sulfide (25 to 40 percent) fractions of the sediment (Kersten and Förstner, 1986). Coarse-grained sediments contain a greater proportion of the total chromium in the non-bioavailable, residual fraction; clayey, organic-rich sediments contain a greater proportion of the total chromium in the more bioavailable organic fraction. More than 70 percent of the chromium in uncontaminated sediments may be associated with the non-bioavailable, residual fraction (Prohic and Kniewald, 1987). The residual chromium is associated primarily with the heavy minerals chromite, chromiferous magnetite, and spinels, as well as with the aluminosilicate lattice of clays (Mayer and Fink, 1980).

6.3.2 Bioavailability and Toxicity in Sediments

Marine and freshwater organisms have evolved efficient mechanisms for bioaccumulating and regulating chromium and other essential trace metals (Simkiss and Taylor, 1989). Concentrations of essential metals (including arsenic, chromium, copper, nickel, and zinc) in tissues of aquatic organisms are regulated at relatively constant values over a wide range of concentrations in the ambient media or food (Chapman et al., 1996). Chromium (III) compounds, because of their low aqueous solubilities, have a low bioavailability to freshwater and marine organisms. Chromium bioaccumulated by marine animals tends to be sequestered in insoluble granules and is not bioavailable to predators of the marine animals (Nott and Nicolaidou, 1996).

Hexavalent chromium is moderately toxic, and trivalent chromium, because of its low aqueous solubility, is practically non-toxic to aquatic organisms. The U.S. EPA acute and chronic marine water quality criteria for chromate are 1,100 $\mu\text{g/L}$ and 50 $\mu\text{g/L}$, respectively (Table 6-1). Marine sediment ERL and ERM values for chromium are 81 $\mu\text{g/g}$ and 370 $\mu\text{g/g}$, respectively.

6.4 Copper

6.4.1 Predominant Forms in Sediment

Concentrations of copper in uncontaminated estuarine and marine sediments are in the range of 10 to 50 $\mu\text{g/g}$ dry wt (Salomons and Förstner, 1984) (Table 6-1). The “high” concentration of copper in marine sediments is 42 $\mu\text{g/g}$ (Daskalakis and O’Connor, 1995). Approximately 25 percent of coastal sediments monitored as part of U.S. monitoring programs contain concentrations of copper equal to or higher than the high value.

Much of the copper in sediments containing low concentrations of organic matter is in the residual fraction associated with the silicate lattice of clays (Chester et al., 1988). In sediments containing high concentrations of organic matter, copper is associated primarily with the organic/sulfide fraction or with extractable organic matter (Luoma, 1985) (Table 2-3). Much of the remainder of the copper in oxidized sediments is associated with the reducible iron and manganese oxides (Prohic and Kniewald, 1987). In anoxic sediments, copper may undergo a variety of reactions with different inorganic and organic sulfur species to form a variety of soluble and insoluble complexes (Shea and Helz, 1988). Polysulfide complexes with cuprous copper (I) are soluble. Thus, the dominant form of copper in solution in the pore water of anoxic sediment layers is $\text{CuS}(\text{S}_5)^{-2}$. The dominant forms of copper in the solid phase of sediment include chalcocite (Cu_2S), covellite (CuS), and possibly chalcopyrite (CuFeS_2) (Shea and Helz, 1988). These sulfides have a low mobility and bioavailability.

6.4.2 Bioavailability and Toxicity in Sediments

Copper is an essential trace nutrient and is bioaccumulated by aquatic organisms primarily from the water. The most bioavailable forms of copper to aquatic organisms are the inorganic hydroxide complexes [CuOH^+ , $\text{Cu}(\text{OH})_2$, $\text{Cu}(\text{OH})_3$, and $\text{Cu}_2(\text{OH})_2$] (Simkiss and Taylor, 1989). The free ion (Cu^{+2}) also is bioavailable (Phinney and Bruland, 1994). Most organic complexes of copper are bioaccumulated inefficiently. Aquatic organisms regulate concentrations of copper in their tissues within a narrow, species-specific range and net accumulation to higher than natural concentrations occurs only when concentrations of bioavailable forms of copper in water or sediments greatly exceed natural levels. Water is the main source of copper in tissues of aquatic organisms (Ettanjani et al., 1992). Copper does not biomagnify in aquatic food webs (Schafer et al., 1982).

Dissolved, reactive copper is toxic to aquatic plants and animals. Free ionic copper at concentrations as low as 0.3 $\mu\text{g/L}$ decreases primary production in several species of oceanic phytoplankton (Brand et al., 1986). However, most of the copper in sea water is complexed with organic matter or in less toxic, bioavailable forms. The U.S. EPA acute and chronic marine water quality criteria for copper are 4.8 $\mu\text{g/L}$ and 3.1 $\mu\text{g/L}$, respectively (Table 6-1). The ERL and ERM for copper in marine sediments are 34 and 270 $\mu\text{g/g}$, respectively (Long et al., 1995).

6.5 Lead

6.5.1 Predominant Forms in Sediment

Concentrations of lead in uncontaminated estuarine and nearshore marine sediments generally fall in the range of 5 to 30 $\mu\text{g/kg}$ dry wt (Salomons and Förstner, 1984) (Table 6-1). Freshwater sediments may contain lower concentrations. The “high” concentration of lead in marine sediments is 45 $\mu\text{g/kg}$ (Daskalakis and O’Connor, 1995). Most of the lead in sediments is associated with fine-grain sediment particles (Krumgalz et al., 1992).

Residual lead (part of the mineral matrix of sediment particles) in uncontaminated sediments, which may represent up to 80 percent of the total lead, is associated primarily with aluminosilicates, sulfide minerals, and barite (Loring, 1982). This residual lead is immobile and not bioavailable. The non-residual lead in oxidized surficial sediments appears to be associated primarily with reducible iron and manganese oxide coatings on clay particles (Luoma and Bryan, 1981) (Table 2-3), as indicated by the strong positive correlation between concentrations of aluminum (from aluminosilicate clay particles) and lead in sediments (Schropp et al., 1990).

In anoxic (oxygen-depleted) sediments, the most stable valence state of lead is the +2 state (Harada and Tsunogai, 1988). Divalent lead (Pb^{+2}) reacts with inorganic sulfide in sediment to form highly insoluble lead sulfide (PbS) (Kersten and Förstner, 1986). However, in highly reducing sediments with an Eh of less than about -0.4 volts, lead may form bisulfide complexes with sulfur. These bisulfide complexes are slightly soluble and the dissolved lead may be mixed up into the water column by sediment disturbance (Shea and MacCrehan, 1988). Most of the lead in oxidized and anoxic sediments is in insoluble and non-bioavailable forms.

6.5.2 Bioavailability and Toxicity in Sediments

Marine deposit-feeding clams and polychaete worms are able to bioaccumulate lead from oxidized sediments (Luoma, 1985). The bioavailability of lead to sediment-associated animals is proportional to the lead/iron concentration ratio in weak acid extracts of the sediment, indicating that the lead absorbed to iron oxide coatings on sediment particles is not bioavailable. In moderately hypoxic or anoxic sediments, most of the lead is precipitated as lead sulfide and is not bioavailable (Bourgoin et al., 1991). Lead is biodepleted in marine food chains relative to calcium, which behaves similarly to lead in the environment (Smith et al., 1990), meaning that it does not biomagnify.

Inorganic lead is moderately toxic to freshwater and marine organisms. U.S. EPA acute and chronic water criteria for inorganic lead for protection of marine life are $220 \mu\text{g/L}$ and $8.5 \mu\text{g/L}$, respectively (Table 6-1). The ERL and ERM concentrations in marine sediments are $46.7 \mu\text{g/g}$ and $218 \mu\text{g/g}$, respectively.

6.6 Mercury

6.6.1 Predominant Forms in Sediment

Concentrations of total mercury in uncontaminated estuarine and marine sediments generally are $0.2 \mu\text{g/g}$ dry wt or lower (Salomons and Förstner, 1984) (Table 6-1), except in areas of natural mercury-containing deposits, such as the East Pacific Rise and the Mid-Atlantic Ridge (Jonasson and Boyle, 1972). The “high” concentration of mercury in coastal sediments is $0.22 \mu\text{g/g}$ (Daskalakis and O’Connor, 1995).

Mercury may occur in three valence states in water and sediments: zero (elemental mercury), +1 (mercurous compounds), and +2 (mercuric compounds) (Moore and Ramamoorthy, 1984). The +2 valence state is the most common in well-oxygenated and hypoxic aquatic environments. Mercury (II) is reduced to elemental mercury, mercuric sulfide, and methylmercury in anoxic sediments (Weber et al., 1998).

Most of the labile (non-residual) mercury in sediments is complexed with particulate and dissolved organic matter in the sediments and not with clay particles or iron oxide coatings on clay particles (Table 2-3). Inorganic and organic mercury salts form very strong and stable complexes with organic ligands in

water and sediment (Moore and Ramamoorthy, 1984). These organic complexes have a low bioavailability to aquatic organisms.

Most mercury methylation takes place in hypoxic or anoxic sediment layers (Gagnon et al., 1996). Mercury methylation is performed primarily by sediment-dwelling, sulfate-reducing bacteria. Under certain conditions, volatile dimethylmercury also is formed (Weber et al., 1998). It may diffuse through the sediment layers into the overlying water column from which it evaporates into the atmosphere. Elemental mercury, also produced by sulfate-reducing bacteria, is slightly volatile and may be lost rapidly from sediments to the atmosphere (Nakamura et al., 1990). In oxidized sediment layers, methylmercury is demethylated to produce inorganic divalent mercury. Because of rapid interconversions of inorganic and organic mercury in oxidized and reduced layers of freshwater and marine sediments, methylmercury rarely represents more than 1 percent of the total mercury in sediments (Berman and Bartha, 1986). Dissolved methylmercury may represent up to about 30 percent of the total dissolved mercury in sediment pore water, but less than 1 percent of the methylmercury adsorbed to sediment particles in the anoxic layers of sediments (Gagnon et al., 1996). Although much of the dissolved methylmercury in sediment pore water is actually complexed to dissolved organic matter, particularly fulvic acids, it should be considered potentially bioavailable to sediment-dwelling organisms. The main pathway for movement of methylmercury from anoxic pore water into the overlying water column is through bioaccumulation by sediment-dwelling animals that are part of the aquatic food web.

High concentrations of sulfide in sediments may inhibit methylmercury formation (Berman and Bartha, 1986). This is thought to be due to formation of extremely insoluble mercuric sulfide (solubility product $10^{-52.4}$). Mercuric sulfide tends to be quite stable and non-bioavailable in hypoxic and anoxic sediments. However, if sulfide concentrations are very high, more soluble disulfide (HgS_2^{-2}) or polysulfide complexes may be formed. These sulfides are more soluble than HgS (Lu and Chen, 1977).

6.6.2 Bioavailability and Toxicity in Sediments

Because of their high affinity for dissolved and particulate organic matter, both inorganic and organic mercury readily complex with organic matter in water and sediments. Mercury bound to organic particles has a low bioavailability to freshwater and marine organisms (Jenne and Luoma, 1977). Methylmercury is more readily bioaccumulated than inorganic mercury (Phillips and Buhler, 1978). This probably is a result of the much slower release of bioaccumulated organic than inorganic mercury by aquatic animals (Thompson, 1990).

Quantitatively, the most important sources of mercury, particularly methylmercury, in the tissues of aquatic animals are probably from ingestion of mercury-contaminated sediments and food. Methylmercury in the tissues of aquatic animals is derived from microbial methylation of inorganic mercury in hypoxic and anoxic layers in the water column and sediments (Rolfhus and Fitzgerald, 1995; Gagnon et al., 1996). The dominant form of mercury in the tissues of most freshwater and marine animals is methylmercury. The concentration of organo-mercury tends to increase with increasing trophic level in aquatic food webs, indicating that organic mercury compounds can be biomagnified in aquatic food webs (Schafer et al., 1982). Very high concentrations of total mercury may be present in the livers of fish-eating marine birds and mammals (Neff, 1997b).

Mercury as the reactive, free inorganic ion and as various organo-mercury compounds in solution is one of the most toxic metals to marine organisms. Acutely toxic concentrations of inorganic mercury in solution are in the range of 3 to 1,000 $\mu\text{g/L}$. However, mercury that is complexed with dissolved or particulate organic matter in the water is not readily bioavailable and has a low aquatic toxicity. The U.S. EPA chronic marine water quality criterion for mercury (II) is 1.106 $\mu\text{g/L}$; the chronic value for

methylmercury is 0.025 µg/L (Table 6-1). However, methylmercury rarely represents more than 10 percent of total mercury in oxygenated surface waters (Mason and Fitzgerald, 1993). Therefore, the chronic value for this form of mercury rarely is exceeded in surface waters. The sediment screening levels for total mercury are 0.15 µg/g ERL and 0.71 µg/g ERM.

6.7 Nickel

6.7.1 Predominant Forms in Sediment

Nickel often is relatively abundant in soils and sediments. Uncontaminated estuarine and marine sediments usually contain 50 µg/g dry wt or less of nickel, the concentration often being positively correlated with the clay content of the sediments (Bowen, 1979) (Table 6-1). The “high” concentration of nickel in sediments from coastal areas of the United States is 42 µg/g (Daskalakis and O’Connor, 1995). However, much higher concentrations of nickel are reported frequently in apparently uncontaminated sediments (Breckenridge and Crockett, 1995). Some soils and sediments, particularly of deep-sea origin, may contain up to 1,000 µg/g nickel (Loring and Asmund, 1996). Similarly, igneous rocks contain 2 to 3,600 ppm nickel (Adriano, 1986), and volcanic minerals may contain high nickel concentrations.

In oxidized sediments, much of the potentially bioavailable nickel is complexed to iron and manganese oxides (Luther et al., 1986) (Table 2-3). Nickel forms weak coordination complexes with oxygen donors such as carboxylate, hydroxyl, and other oxy-ligands (e.g., humic and fulvic acids, clays, and metal oxides) (Wood, 1987). It also becomes tightly bound to anionic groups of bacterial polysaccharides (Wood, 1987). Nickel forms stable, insoluble complexes with surfides and organic thiols in anoxic sediment layers (Wood, 1987). However, most of the nickel (often more than 90 percent) in relatively uncontaminated sediments is in the residual fraction, associated primarily with oxide minerals, such as magnetite, spinels, and silicates (Loring, 1982). Thus, the bioavailability of nickel in sediments usually is low.

6.7.2 Bioavailability and Toxicity in Sediments

Like other essential metals, nickel concentrations in the tissues of aquatic organisms do not covary with nickel concentrations in the ambient water, sediments, and prey items. Of the dominant forms of nickel in sediments and sediment pore water [Ni^{+2} , $\text{Ni}(\text{OH})_2$, and NiS], only nickel ion is readily bioavailable (Förstner and Wittmann, 1981). However, nickel sulfide is the most soluble of the common metal sulfides and readily dissolves when the oxygen concentration in sediment increases. Similarly, nickel weakly complexed to organic matter in surface sediments readily exchanges with divalent cations in the water, releasing bioavailable nickel ion to the overlying water column (Morse, 1995). The hydroxide and sulfide are insoluble. Nickel in soils generally is not bioavailable to earthworms (Sample et al., 1998).

Inorganic nickel has a relatively low toxicity to aquatic organisms. The U.S. EPA marine acute and chronic water quality criteria for nickel are 75 µg/L and 8.3 µg/L, respectively (Table 6-1). ERL and ERM values for nickel in marine sediments are 20.9 µg/g and 51.6 µg/g, respectively (Long et al., 1995). These screening values often are exceeded (usually without adverse effects in benthic organisms) as a result of the high abundance of residual nickel in several crustal rocks and minerals.

6.8 Tin

6.8.1 Predominant Forms in Sediment

The concentration of inorganic tin in uncontaminated sediments is about 2 µg/g dry wt. Although inorganic tin compounds may be moderately toxic to aquatic organisms, contamination of aquatic ecosystems with inorganic tin is rarely perceived as a problem, except possibly near some metal smelting and mining operations (Skei et al., 1972). However, various organotin compounds, some of which are extremely toxic to aquatic organisms, are used for a variety of commercial purposes that favor their entry into the marine environment. Most organotins contain tetravalent tin covalently bonded to one to four organic substituents (Müller et al., 1989). Tripropyl-, tributyl-, and triphenyl-tins are extremely effective biocides that are used as wood preservatives, antifoulants for boat hulls and other submerged structures, and disinfectants and slimicides for cooling and paper mill waters (Snoeij et al., 1987). Although organotins do not adsorb strongly to particles, they do tend to accumulate in sediments in the vicinity of major sources in the water column (e.g., marinas and ship yards), though their concentrations rarely are as high as those of inorganic tin.

Tributyltin (the most common organotin in antifouling coatings) is present in aerobic sediment primarily as tributyltin chloride, tributyltin hydroxide, and tributyltin carbamate (Eng et al., 1986). In anaerobic sediment, the dominant chemical forms appear to be the sulfide, hydroxide, and carbonate. Tributyltins undergo sequential de-alkylation in sediments to yield dibutyltin, monobutyltin, and finally inorganic tin (Maguire and Tkacz, 1985). The degradation half-life of tributyltin in oxidized marine sediments is approximately 162 days (Stang and Seligman, 1986). Biodegradation of tributyltin in hypoxic or anoxic sediments is negligible.

6.8.2 Bioavailability and Toxicity in Sediments

Organotins in water, sediments, and tissues of aquatic organisms are relatively bioavailable (Laughlin and French, 1988). They also are highly toxic to aquatic organisms (Langston et al., 1990).

Concentrations as low as 1-2 ng/L (parts per billion) of dissolved tributyltin causes severe reproductive and developmental effects in freshwater and marine invertebrates. These concentrations are observed in the water of marinas and ports where vessels are protected with tributyltin-based paints from biofouling organisms. Because of their high toxicity, tributyltin antifouling paints recently were banned for most marine and freshwater uses in the United States and Europe.

6.9 Zinc

6.9.1 Predominant Forms in Sediment

Concentrations of zinc in uncontaminated sediments vary widely. Coarse-grained sandy sediments may contain as little as 1.2 µg/g dry wt zinc; clay sediments may contain more than 100 µg/g total zinc (Larsen and Gaudette, 1995) (Table 6-1). The “high” concentration of zinc in U.S. coastal sediments is 135 µg/g (Daskalakis and O’Connor, 1995).

Most of the zinc in sediments is residual, rendering it non-bioavailable. The residual zinc is associated with the mineral lattice of clays and with a variety of heavy minerals, including chromite, ilmenite, and magnetite (Loring, 1982). Sphalerite (ZnS) and zincite (ZnO) are important carriers of residual zinc in some sediments. The nonresidual zinc in many oxidized sediments is associated primarily with the reducible iron and manganese oxide fractions. In reducing sediments, much of the zinc is associated with

the organic/sulfide fraction (Rosental et al., 1986) (Table 2-3). During transitions of oxidation/reduction potential in sediments, zinc may be released in soluble form into sediment pore water, from which it diffuses into the overlying water column. The total flux of zinc from sediments into the waters of the whole of southern San Francisco Bay is approximately 298 kg/day (Wood et al., 1995).

6.9.2 Bioavailability and Toxicity in Sediments

Zinc is an essential micronutrient in all aquatic organisms, being a cofactor in several enzymes. Most aquatic species have efficient mechanisms for bioaccumulating zinc, and some species store zinc in non-toxic forms in their tissues. Freshwater and marine organisms accumulate zinc from water, food, and sediments. Sediment-dwelling aquatic invertebrates can accumulate zinc adsorbed to iron oxides in oxidized sediments (Harvey and Louma, 1985). Much of the zinc in tissues of aquatic organisms is sequestered in phosphate granules and is not bioavailable to predators (Nott and Nicolaidou, 1993). Zinc is not biomagnified in aquatic food webs.

The toxic species of zinc is the free ion, which represents only a small fraction of the total zinc in natural water and sediment pore water. Acutely lethal concentrations of total zinc in solution usually are in the range of 100 to 50,000 µg/L. Sublethal responses are observed, particularly in aquatic plants, at much lower concentrations. Invertebrates and plants seem to be more sensitive than fish and higher animals to zinc poisoning. The U.S. EPA acute and chronic water quality criteria for zinc are 95 µg/L and 86 µg/L, respectively (Table 6-1). The ERL and ERM for zinc in marine sediments are 150 µg/g and 410 µg/g, respectively, reflecting the relatively low toxicity of sediment-bound zinc (Long et al., 1995).

7.0 SUMMARY OF SELECTED CASE STUDIES

Bioavailability adjustments have been incorporated into human health risk assessments for several sites having metals contamination. The number of such sites continues to grow as the concept of bioavailability is better understood and gains acceptance among the regulatory community. Bioavailability studies have been used both at sites where U.S. EPA is the lead regulatory agency (Regions III, VII, VIII, IX, and X) and at sites where the state agency has the lead (Oklahoma, Michigan, California, Illinois, Wisconsin, and New Jersey). Bioavailability adjustments have been supported by in vivo animal studies, in vitro testing, environmental health studies, mineral speciation, or some combination of these methods. To date, most bioavailability adjustments have been made for the oral route of exposure. Only one case study was identified for dermal bioavailability, and none were identified for the inhalation pathway. Bioavailability adjustments have been made for arsenic, lead, mercury, and cadmium; however, the majority of adjustments have been for lead and arsenic associated with mining and smelting activities.

Results of several case studies are presented in Table 7-1. Most of the case studies presented here illustrate decreased bioavailability compared to the default assumptions and thus increased cleanup levels; however, it should be noted that in some cases (particularly for lead, where the default assumption is 30 percent absolute bioavailability from soil) bioavailability studies can support the default assumption or even demonstrate higher bioavailability than the default. One such example in Table 7-1 is the Palmerton, PA site, where swine studies supported the default bioavailability value of 30 percent.

Among the case studies presented in Table 7-1, the National Zinc Company NPL Site in Bartlesville, OK illustrates several factors that are important in getting a bioavailability study accepted. In this case study, the regulators and other stakeholders were involved from the beginning. A detailed work plan including protocols for the bioavailability studies was prepared. Protocols were developed with input from toxicologists with training in pharmacokinetics to select appropriate animal models and testing endpoints. These protocols followed GLP Standards and were peer reviewed by an outside toxicologist brought in by the stakeholders. Also, the regulators and stakeholders were given the opportunity to review the results prior to making final interpretation. The bioavailability studies for this site supported RAFs of 0.25 for arsenic, 0.33 for cadmium, and 0.20 (vs. default of 0.30) for lead. Using these adjustments for bioavailability, the Oklahoma Department of Environmental Quality (DEQ) accepted a threefold increase in cleanup levels for arsenic and cadmium (from 20 to 60 ppm for arsenic and from 30 to 100 ppm for cadmium) and almost a twofold increase in the cleanup level for lead (from 500 to 925 ppm). In this case, the process from drafting the work plan to draft remedial investigation report for public comment required only seven months. The costs related to the bioavailability studies (work plan development and laboratory testing) were approximately \$200,000; however, the increased cleanup goals reduced remediation costs by approximately \$40 million.

Table 7-1. Selected Case Studies for Bioavailability Adjustments

Site	Contaminant	Test	Bioavailability Test Results	Cleanup Level	Regulatory Agency
National Zinc Co. NPL Site, Bartlesville, OK	Lead	In vivo – rat; and speciation	40% (20% absolute)	925 mg/kg	Oklahoma DEQ
	Cadmium	In vivo – rat; and speciation	33%	100 mg/kg	
	Arsenic	In vitro (PBET); and speciation	25%	60 mg/kg	
Butte, MT	Lead	In vivo – rat	24% (12% absolute)	1,200 mg/kg	U.S. EPA Region VIII
Palmerton, PA	Lead	In vivo – swine, Monte Carlo analysis	30% absolute (same as default)	650 mg/kg	U.S. EPA Region III
Anaconda, MT	Arsenic (soil)	In vivo – monkey	18.3%	250 ppm	U.S. EPA Region VIII
	Arsenic (dust)	In vivo – monkey	25.8%		
Rushton/North Tacoma, WA Off-Site	Arsenic (soil)	None – Regulators accepted adjustment	80%	230 ppm	U.S. EPA Region X
Oak Ridge National Laboratory, TN	Mercury	In vivo, in vitro, speciation	10%	400 ppm	U.S. EPA Region IV
Carson River, NV	Mercury (insoluble 90%, soluble 10%)	Speciation	(20% for insoluble; 100% for soluble) 30% overall	80 ppm	U.S. EPA Region IX
Crego Park, MI	Arsenic	In vitro (PBET) and speciation	10%	68 ppm (from 6.8 ppm)	Michigan DEQ
Almaden Quicksilver County Park, Los Gatos, CA	Mercury	In vitro and speciation	30%	300 to 500 ppm for various areas in park	Cal-EPA DTSC
Union Pacific Railroad Yard, Sacramento, CA	Arsenic	In vivo – swine	0-1% absorption from slag vs. 59% absorption of soluble control	No cleanup required (slag up to 1,800 ppm As)	Cal-EPA DTSC
Hudson Co., NJ	Chromium	In vitro extraction (ASTM method 3987)	Endpoint allergic contact dermatitis	State has recommended test but no results yet	NJDEP

Cal-EPA = California Environmental Protection Agency.

DEQ = Department of Environmental Quality.

DTSC = Department of Toxic Substances Control.

NJDEP = New Jersey Department of Environmental Protection.

PBET = Physiologically Based Extraction Test.

8.0 REFERENCES

- Adriano, D.C. 1986. *Trace Elements in the Terrestrial Environment*. Springer-Verlag, New York, NY.
- Agency for Toxic Substances and Disease Registry. 1993a. *Toxicological Profile for Arsenic*. Prepared by Life Systems, Inc., for U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Agency for Toxic Substances and Disease Registry. 1993b. *Toxicological Profile for Lead*. Prepared by Clement International Corporation, for U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Agency for Toxic Substances and Disease Registry. 1997a. *Toxicological Profile for Cadmium* (draft). U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Agency for Toxic Substances and Disease Registry. 1997b. *Toxicological Profile for Mercury* (draft). U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Agency for Toxic Substances and Disease Registry. 1997c. *Toxicological Profile for Nickel*. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Agency for Toxic Substances and Disease Registry. 1998. *Toxicological Profile for Chromium* (draft). Prepared by Sciences International, Inc., under contract to Research Triangle Institute, under Contract No. 205-93-0606, for U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- Allen, H.E., G. Fu, and B. Deng. 1993. "Analysis of Acid-Volatile Sulfide (AVS) and Simultaneously Extracted Metals (SEM) for Estimation of Potential Toxicity in Aquatic Sediments." *Environ. Toxicol. Chem.*, 12: 1441-1453.
- American Society for Testing and Materials. 1995. *Standard Guide for Risk-Based Corrective Action Applied at Petroleum Release Sites (E 1739-95)*. West Conshohocken, PA.
- American Society for Testing and Materials. 1998. *Standard Provisional Guide for Risk-Based Corrective Action (PS 104-98)*. West Conshohocken, PA.
- Ankley, G.T. 1996. "Evaluation of Metal/Acid-Volatile Sulfide Relationships in the Prediction of Metal Bioaccumulation by Benthic Macroinvertebrates." *Environ. Toxicol. Chem.*, 15: 2138-2146.
- Ankley, G.T., D.M. Di Toro, D.J. Hansen, and W.J. Berry. 1996. "Technical Basis and Proposal for Deriving Sediment Quality Criteria for Metals." *Environ. Toxicol. Chem.*, 15: 2056-2066.
- ASTM, see American Society for Testing and Materials.
- ATSDR, see Agency for Toxic Substances and Disease Registry.
- Berman, M. and R. Bartha. 1986. "Control of the Methylation Process in a Mercury-Polluted Aquatic Sediment." *Environ. Pollut. (Ser. B)*, 11: 41-53.

- Berry, W.J., M.G. Cantwell, P.A. Edwards, J.R. Serbst, and D.J. Hansen. 1999. "Predicting Toxicity of Sediments Spiked with Silver." *Environ. Toxicol. Chem.*, 18: 40-55.
- Bechtel Jacobs Company, LLC. 1998. *Biota Sediment Accumulation Factors for Invertebrates: Review and Recommendations for the Oak Ridge Reservation*. Prepared for the U.S. DOE Office of Environmental Management, Code EW20. Contract No. DE-AC05-98OR22700. BJC/OR-112. August.
- BJC, see Bechtel Jacobs Company, LLC.
- Boulègue, J. 1983. "Trace metals (Fe, Cu, Zn, Cd) in Anoxic Environments." In: C.S. Wong, E. Boyle, K.W. Bruland, J.D. Burton, and E.D. Goldberg (Eds.), *Trace Metals in Sea Water*, pp. 563-578. Plenum Press, New York, NY.
- Bourgoin, B.P., M.J. Risk, R.D. Evans, and R.J. Cornett. 1991. "Relationships Between the Partitioning of Lead in Sediments and its Accumulation in the Marine Mussel, *Mytilus edulis* Near a Lead Smelter." *Wat. Air Soil Pollut.*, 57-58: 377-386.
- Bowen, H.J.M. 1979. *Environmental Chemistry of the Elements*. Academic Press, London.
- Brand, L.E., W.G. Sunda, and R.R.L. Guillard. 1986. "Reduction of Marine Phytoplankton Reproduction Rates by Copper and Cadmium." *J. Exp. Mar. Biol. Ecol.*, 96: 225-250.
- Breckenridge, R.P. and A.B. Crockett. 1995. *Determination of Background Concentrations of Inorganics in Soils and Sediments at Hazardous Waste Sites*. EPA/540/5-96/500. Prepared for U.S. Environmental Protection Agency, Office of Research and Development, Washington, DC.
- Brown, B. and J. Neff. 1993. *Bioavailability of Sediment-Bound Contaminants to Marine Organisms*. Report PNL-8761 UC-0000. Prepared by Battelle Marine Sciences Laboratory for the National Ocean Pollution Program Office, NOAA.
- Bryan, G.W. and W.J. Langston. 1992. "Bioavailability, Accumulation and Effects of Heavy Metals in Sediments with Special Reference to United Kingdom Estuaries: A Review." *Environ. Pollut.*, 76: 89-131.
- Camp, Dresser and McKee, Inc. 1992. *Risk Assessment, Almaden Quicksilver County Park, Volume II—Appendices*. Final report prepared for Santa Clara County Parks and Recreation Department, Los Gatos, CA.
- Campbell, P.G.C., A.G. Lewis, P.M. Chapman, A.A. Crowder, W.K. Fletcher, B. Imber, S.N. Luoma, P.M. Stokes, and M. Winfrey. 1988. *Biologically Available Metals in Sediments*. NRCC No. 27694. National Research Council of Canada, Ottawa, Canada. 298 pp.
- Canli, M. and R.W. Furness. 1995. "Mercury and Cadmium Uptake from Seawater and from Food by the Norway Lobster *Nephrops norvegicus*." *Environ. Toxicol. Chem.*, 14: 819-828.
- Casteel, S.W., R.P. Cowart, C.P. Weis, G.M. Henningsen, E. Hoffman, W.J. Brattin, R.E. Guzman, M.F. Starost, J.T. Payne, S.L. Stockham, S.V. Becker, J.W. Drexler, and J.R. Turk. 1997a. "Bioavailability of Lead to Juvenile Swine Dosed with Soil from the Smuggler Mountain NPL Site of Aspen, Colorado." *Fund. Appl. Toxicol.*, 36: 177-187.

- Casteel, S.W., L.D. Brown, M.E. Dunsmore, C.P. Weis, G.M. Henningsen, E. Hoffman, W.J. Brattin, and T.L. Hammon. 1997b. *Relative Bioavailability of Arsenic in Mining Wastes*. Document Control No. 4500-88-AORH. Prepared for U.S. EPA Region VIII, Denver, CO.
- CDM, see Camp, Dresser and McKee, Inc.
- Chapman, P.M., H. E. Allen, K. Godtfredsen, and M.N. Z'Graggen. 1996. "Evaluation of Bioaccumulation Factors in Regulating Metals." *Environ. Sci. Technol.*, 30: 448A-452A.
- Chester, R., A. Thomas, F.J. Lin, A.S. Basaham, and G. Jacinto. 1988. "The Solid State Speciation of Copper in Surface Water Particles and Oceanic Sediments." *Mar. Chem.*, 24: 261-292.
- Daskalakis, K.D. and T.P. O'Connor. 1995. "Distribution of Chemical Concentrations in U.S. Coastal and Estuarine Sediment." *Mar. Environ. Res.*, 40: 381-398.
- De Vitre, R., N. Belzile, and A. Tessier. 1991. "Speciation and Adsorption of Arsenic on Diagenic Iron Oxyhydroxides." *Limnol. Oceanogr.*, 36: 1480-1485.
- Dieter, M.P., H.B. Matthews, R.A. Jeffcoat, and R.F. Moseman. 1993. *Comparison of Lead Bioavailability in F344 Rats Fed Lead Acetate, Lead Oxide, Lead Sulfide, or Lead Ore*.
- DiToro, D.M., J.D. Mahony, D.J. Hansen, K.J. Scott, M.B. Hicks, S.M. Mayr, and M.S. Redmond. 1990. "Toxicity of Cadmium in Sediments: the Role of Acid Volatile Sulfide." *Environ. Toxicol. Chem.*, 9: 1487-1502.
- Eisler, R. 1985. *Cadmium Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review*. Biological Report 85(1-2). Contaminant Hazard Reviews. Report No. 2. U.S. Dept. of the Interior, Fish and Wildlife Service, Washington, DC.
- Eng, G., O. Bathersfield, and L. May. 1986. "Mössbauer Studies of the Speciation of Tributyltin Compounds in Seawater and Sediment Samples." *Water Air Soil Pollut.*, 27: 191-197.
- Ettanjani, H. C. Amiard-Triquet, and J.-C. Amiard. 1992. "Etude Expérimentale du Transfert de Deux Éléments Traces (Ag, Cu) dans une Chaîne Trophique Marine: Eau - Particules (Sédiment Natural, Microalgue) - Mollusques Filtreurs (*Crassostrea gigas* Thunberg)." *Wat. Air Soil Pollut.*, 65: 215-236.
- Förstner, U. and G.T.W. Wittmann. 1981. *Metal Pollution in the Aquatic Environment*. Second Revised Edition. Springer-Verlag, Berlin.
- Freeman, G.B., J.D. Johnson, J.M. Killinger, S.C. Liao, P.I. Feder, A.O. Davis, M.V. Ruby, R.L. Chaney, S.C. Lovre, and P.D. Bergstrom. 1992. "Relative Bioavailability of Lead from Mining Waste Soil in Rats." *Fund. Appl. Toxicol.*, 19: 388-398.
- Freeman, G.B., J.D. Johnson, J.M. Killinger, S.C. Liao, A.O. Davis, M.V. Ruby, R.L. Chaney, S.C. Lovre, and P.D. Bergstrom. 1993. "Bioavailability of Arsenic in Soil Impacted by Smelter Activities Following Oral Administration in Rabbits." *Fundam. Appl. Toxicol.*, 21: 83-88.
- Freeman, G.B., J.D. Johnson, S.C. Liao, P.I. Feder, A.O. Davis, M.V. Ruby, R.A. Schoof, R.L. Chaney, and P.D. Bergstrom. 1994. "Absolute Bioavailability of Lead Acetate and Mining Waste Lead in Rats." *Toxicology*, 91: 151-163.

- Freeman, G.B., R.A. Schoof, M.V. Ruby, A.O. Davis, J.A. Dill, S.C. Liao, C.A. Lapin, and P.D. Bergstrom. 1995. "Bioavailability of Arsenic in Soil and House Dust Impacted by Smelter Activities Following Oral Administration in Cynomolgus Monkeys." *Fundam. Appl. Toxicol.*, 28: 215-222.
- Freeman, G.B., J.A. Dill, J.D. Johnson, P.J. Kurtz, F. Parham, and H.B. Matthews. 1996a. "Comparative Absorption of Lead from Contaminated Soil and Lead Salts by Weanling Fischer 344 Rats." *Fundam. Appl. Toxicol.*, 33: 109-119.
- Freeman, G.B., J.A. Dill, and N.J. Trigg. 1996b. *Determination of the bioavailability of soluble arsenic and arsenic in slag following oral administration in microswine*. Prepared for Dames & Moore, Sacramento, California, by Battelle Columbus Operations, Columbus, OH.
- Gagnon, C., É. Pelletier, A. Mucci, and W.F. Fitzgerald. 1996. "Diagenic Behavior of Methylmercury in Organic-Rich Coastal Sediments." *Limnol. Oceanogr.*, 41: 428-434.
- Gargas, M.L., R.L. Norton, M.A. Harris, D.J. Paustenbach, and B.L. Finley. 1994. "Urinary Excretion of Chromium Following Ingestion of Chromite-Ore Processing Residues in Humans: Implications for Biomonitoring." *Risk. Anal.*, 14(6): 1019-24.
- Goldberg, E.D. 1954. "Marine Geochemistry: Chemical Scavengers of the Sea." *J. Geol.*, 62: 249-266.
- Griffin, S.R., R. Rubenstein, S. Irene, C. DeRosa, and H. Choudhury. 1990. "Bioavailability in Rats of Metals Adsorbed to Soils." U.S. Environmental Protection Agency, Washington, DC, Hazleton Laboratories, America, Inc. Poster presented at the Society of Toxicology 29th Annual Meeting, Miami Beach, FL. February 12-16, 1990. Poster paper no. 623.
- Groen, K., H. Vaessen, J.J.G. Kliest, J.L.M. deBoer, T.V. Ooik, A. Timmerman, and F.F. Vlug. 1994. "Bioavailability of Inorganic Arsenic from Bog Ore-Containing Soil in the Dog." *Environ. Health Perspect.*, 102(2): 182-184.
- Harada, K. and S. Tsunogai. 1988. "Is Lead Soluble at the Surface of Sediments in Biologically Productive Seas?" *Cont. Shelf Res.*, 8: 387-396.
- Hardy, J.T., M.F. Sullivan, E.A. Crecelius, and C.W. Apts. 1984. "Transfer of Cadmium in a Phytoplankton-Oyster-Mouse Food Chain." *Arch Environ. Contam. Toxicol.*, 13: 419-425.
- Harvey, R.W. and S.N. Luoma. 1985. "Separation of Solute and Particulate Vectors of Heavy Metal Uptake in Controlled Suspension-Feeding Experiments with *Macoma balthica*." *Hydrobiol.*, 121: 97-102.
- Hayes, K.F. and S.J. Traina. 1998. "Metal Ion Speciation and its Significance in Ecosystem Health." In: *Soil Chemistry and Ecosystem Health, Special Publication No. 52*. Soil Science Society America, Madison, WI.
- Horowitz, S.B. and B.L. Finley. 1993. "Using Human Sweat to Extract Chromium from Chromite or Processing Residue: Applications to Setting Health-Based Cleanup Levels." *J. Toxicol. Environ. Health*: 585-599.

- Hrudey, S.E., W. Chen, and C.G. Rousseaux. 1996. "Exposure Routes and Bioavailability Factors for Selected Contaminants. I. Arsenic and III. Chromium and Chromium Compounds." In: S.E. Hrudey (Ed.), *Bioavailability in Environmental Risk Assessment*. CRC Press, Boca Raton, FL.
- Jenne, E.A. and S.N. Luoma. 1977. "Forms of Trace Elements in Soils, Sediments, and Associated Waters: an Overview of Their Determination and Biological Availability." In: R.E. Widlung and H. Drucker (Eds.), *Biological Implications of Metals in the Environment*, pp. 110-143. CONF-750929, NTIS, Springfield, VA.
- Jonasson, I.R. and R.W. Boyle. 1972. "Geochemistry of Mercury and Origins of Natural Contamination of the Environment." *CIM Trans.*, 75: 8-15.
- Kersten, M. and U. Förstner. 1986. "Chemical Fractionation of Heavy Metals in Anoxic Estuarine and Coastal Sediments." *Water Sci. Technol.*, 18: 121-130.
- Krumgalz, B.S., G. Fainshtein, and A. Cohen. 1992. "Grain Size Effect on Anthropogenic Trace Metal and Organic Matter Distribution in Marine Sediments." *Sci. Tot. Environ.*, 116: 15-30.
- Langston, W.J., G.W. Bryan, G.R. Burt, and P.E. Gibbs. 1990. "Assessing the Impact of Tin and TBT in Estuaries and Coastal Regions." *Funct. Ecol.*, 4: 433-443.
- Larsen, P.F. and H. Gaudette. 1995. "Spatial and Temporal Aspects of Sedimentary Trace Metal Concentrations in Mid-Coast Maine." *Mar. Pollut. Bull.*, 30: 437-444.
- Laughlin, R.B., Jr. and W. French. 1988. "Concentration Dependence of bis(tributyl)tin Oxide Accumulation in the Mussel, *Mytilus edulis*." *Environ. Toxicol. Chem.*, 7: 1021-1026.
- Lee, F.Y. and J.A. Kittrick. 1984. "Elements Associated with the Cadmium Phase in a Harbor Sediment as Determined with the Electron Beam Microprobe." *J. Environ. Qual.*, 13: 337-340.
- Long, E.R., D.D. Macdonald, S.L. Smith, and F.D. Calder. 1995. "Incidence of Adverse Biological Effects Within Ranges of Chemical Concentrations in Marine and Estuarine Sediments." *Environ. Manage.*, 19(1): 81-97.
- Loring, D.H. 1982. "Geochemical Factors Controlling the Accumulation and Dispersal of Heavy Metals in Bay of Fundy Sediments." *Can. J. Earth Sci.*, 19: 930-944.
- Loring, D.H. and G. Asmund. 1996. "Geochemical Factors Controlling the Accumulation of Major and Trace Elements in Greenland Coastal and Fjord Sediments." *Environ. Geol.*, 28: 2-11.
- Loutit, M., P. Bremer, and J. Aislable. 1988. "The Significance of the Interactions of Chromium and Bacteria in Aquatic Habitats." In: J.O. Nriagu and E. Nieboer (Eds.), *Chromium in the Natural and Human Environments*, pp. 317-334. John Wiley and Sons, New York, NY.
- Lu, J.C.S. and K.Y. Chen. 1977. "Migration of Trace Metals in Interfaces of Seawater and Polluted Surficial Sediments." *Environ. Sci. Technol.*, 11: 174-182.
- Luoma, S.N. 1985. "Biological Availability of Sediment-Bound Trace Metals." In: *La Baie de Seine (Greco-Manche)*, pp. 347-362. IFRAMER, Actes de Colloques No. 4.

- Luoma, S.N. 1989. "Can We Determine the Biological Availability of Sediment-Bound Trace Elements?" *Hydrobiologia*, 176/177: 379-396.
- Luoma, S.N. and G.W. Bryan. 1981. "A Statistical Assessment of the Form of Trace Metals in Oxidized Estuarine Sediments Employing Chemical Extractions." *Sci. Tot. Environ.*, 17: 165-196.
- Luther, G.W. III, Z. Wilk, R.A. Ryans, and L. Meyerson. 1986. "On the Speciation of Metals in the Water Column of a Polluted Estuary." *Mar. Pollut. Bull.*, 17: 535-542.
- Maddaloni, M., P. Goodrum, G. Diamond, W. Manton, C. Blum, N. LoLacono, and J. Graziano. 1998. "Metal-Weighted Bioavailability of Soil-Borne Lead in Adults: Use of Empirical Data Analyzed by Stable Isotope Dilution, and Probabilistic Worker Exposure Scenarios." Poster presented at the Society of Toxicology, 37th Annual Meeting in Seattle, WA.
- Maguire, R.J. and R.T. Tkacz. 1985. "Degradation of tri-*n*-butyltin Species in Water and Sediment from Toronto Harbor." *J. Agric. Food Chem.*, 33: 947-953.
- Mason, R.P. and W.F. Fitzgerald. 1993. "The Distribution and Biogeochemical Cycling of Mercury in the Equatorial Pacific Ocean." *Deep-Sea Res.*, 20: 1897-1924.
- Masscheleyn, P.H., R.D. Delaune, and W.H. Patrick, Jr. 1991. "Arsenic and Selenium Chemistry as Affected by Sediment Redox Potential and pH." *J. Environ. Qual.*, 20: 522-527.
- Mayer, L.M. 1988. "Geochemistry of Chromium in the Oceans." In: J.O. Nriagu and E. Nieboer (Eds.), *Chromium in Natural and Human Environments*, pp. 173-187. John Wiley and Sons, New York, NY.
- Mayer, L.M. and K.F.R. Fink. 1980. "Granulometric Dependence of Chromium Accumulation in Estuarine Sediments in Maine." *Estuar. Cstl. Mar. Sci.*, 11: 491-503.
- Medlin, E.A. 1997. *An In Vitro Method for Estimating the Relative Bioavailability of Lead in Humans*. M.S. Thesis. Department of Geological Sciences, University of Colorado at Boulder.
- Miller, D.D. and B.R. Schriker. 1982. "In Vitro Estimation of Food Iron Bioavailability." In: C. Kies (Ed.), *Nutritional Bioavailability of Iron*. ACS Symposium Series No. 203, American Chemical Society.
- Miller, D.D., B.R. Schriker, R.R. Rasmussen, and D. Van Campen. 1981. "An In Vitro Method for Estimation of Iron Availability from Meals." *Am. J. Clin. Nutr.*, 34: 2248-2256.
- Moore, J.W. and S. Ramamoorthy. 1984. "Heavy Metals in Natural Waters." *Applied Monitoring and Risk Assessment*. Springer Verlag, New York, NY.
- Morse, J.W. 1994. "Interactions of Trace Metals with Authigenic Sulfide Minerals: Implications for Their Bioavailability." *Mar. Chem.*, 46: 1-6.
- Morse, J.W. 1995. "Dynamics of Trace Metal Interactions with Authigenic Sulfide Minerals in Anoxic Sediments." In: H.E. Allen (Ed.), *Metal Contaminated Aquatic Sediments*, pp. 187-199. Ann Arbor Press, Ann Arbor, MI.

- Müller, M.D., L. Renberg, and G. Rippin. 1989. "Tributyltin in the Environment – Sources, Fate, and Determination. An Assessment of Present Status and Research Needs." *Chemosphere*, 18: 2015-2042.
- Nakamura, K., M. Sakamoto, H. Uchiyama, and O. Yagi. 1990. "Organomercurial-Volatilizing Bacteria in the Mercury-Polluted Minamata Bay, Japan." *Appl. Environ. Microbiol.* (Jan. 1990): 304-305.
- National Environmental Policy Institute. 1998. *Bioavailability: Implications for Science/Cleanup Policy*. Bioavailability Policy Project. White Paper.
- Neff, J.M. 1997a. "Ecotoxicology of Arsenic in the Marine Environment: A Review." *Environ. Toxicol. Chem.*, 16: 917-927.
- Neff, J.M. 1997b. *Metals and Organic Chemicals Associated with Oil and Gas Well Produced Water: Bioaccumulation, Fates, and Effects in the Marine Environment*. Report to the Gulf of Mexico Offshore Operators Committee, New Orleans, LA.
- Nelson, A. and P. Donkin. 1985. "Processes of Bioaccumulation: the Importance of Chemical Speciation." *Mar. Pollut. Bull.*, 16: 164-169.
- NEPI, see National Environmental Policy Institute.
- Nielsen, J.B., and O. Andersen. 1990. "Disposition and Retention of Mercuric Chloride in Mice After Oral and Parenteral Administration." *J. Toxicol. Environ. Health*, 30(3): 167-180.
- Newman, M.C. and C.H. Jagoe. 1994. "Ligands and the Bioavailability of Metals in Aquatic Environments." In: J.L. Hamelink, P.F. Landrum, H.L. Bergman, and W.H. Benson (Eds.), *Bioavailability: Physical, Chemical and Biological Interactions*. CRC Press, Inc. Boca Raton, FL.
- Nott, J.A. and A. Nicolaidou. 1993. "Bioreduction of Zinc and Manganese Along a Molluscan Food Chain." *Comp. Biochem. Physiol.*, 104A: 235-238.
- Nott, J.A. and A. Nicolaidou. 1996. "Kinetics of Metals in Molluscan Faecal Pellets and Mineralized Granules, Incubated in Marine Sediments." *J. Exper. Mar. Biol. Ecol.*, 197: 203-218.
- Olazabal, M.A., N.P. Nikolaidis, S.A. Suib, and J.M. Madaraiga. 1997. "Precipitation Equilibria of the Chromium (VI)/Iron (III) System and Spectroscopic Characterization of the Precipitates." *Environ. Sci. Technol.*, 31: 2898-2902.
- Peltonen, L. 1979. "Nickel Sensitivity in the General Population." *Contact Dermatitis*, 5: 27-32.
- Phillips, G.R. and D.R. Buhler. 1978. "The Relative Contributions of Methylmercury from Food and Water to Rainbow Trout (*Salmo gairdneri*) in a Controlled Laboratory Environment." *Trans. Amer. Fish. Soc.*, 107: 853-861.
- Phinney, J.T. and K.W. Bruland. 1994. "Uptake of Lipophilic Organic Cu, Cd, and Pb Complexes in the Coastal Diatom *Thalassiosira weissflogii*." *Environ. Sci. Technol.*, 28: 1781-1790.
- Prohic, E. and G. Kniewald. 1987. "Heavy Metal Distribution in Recent Sediments of the Krka River Estuary: An Example of Sequential Extraction Analysis." *Mar. Chem.*, 22: 279-297.

- PTI Environmental Services. 1994. *Remedial Investigation Report: National Zinc Site Remedial Investigation and Feasibility Study, Volume 1*. Prepared for City of Bartlesville, Cyprus Amax Minerals Company, and Salomon, Inc.
- PTI, see PTI Environmental Services.
- Rahola, T., T. Hattula, A. Korhonen, and J.K. Miettinen. 1973. "Elimination of Free and Protein-Bound Ionic Mercury ($^{203}\text{Hg}^{2+}$) in Man." *Ann. Clin. Res.*, 5: 214-219.
- Revis, N., G. Holdsworth, G. Bingham, A. King, and J. Elmore. 1989. *An Assessment of Health Risk Associated with Mercury in Soil and Sediment from East Fork Poplar Creek, Oak Ridge, Tennessee*. U.S. Department of Energy, Oak Ridge Research Institute, Oak Ridge, TN.
- Revis, N.W., T.R. Osborne, G. Holdsworth, and C. Hadden. 1990. "Mercury in Soil: A Method for Assessing Acceptable Limits." *Arch. Environ. Contam. Toxicol.*, 19: 221-226.
- Rodriguez, R.R., N.T. Basta, S.W. Casteel, and L.W. Pace. 1999. "An In Vitro Gastrointestinal Method to Estimate Bioavailable Arsenic in Contaminated Soils and Solid Media." *Environ. Sci. Technol.*, 33(4): 642-649.
- Rodriguez, V.M., L. Dufour, L. Carrizales, F. Diaz-Barriga, and M.E. Jiminez-Capdeville. 1998. "Effects of Oral Exposure to Mining Waste on *In Vivo* Dopamine Release from Rat Striatum." *Environ. Health Perspect.*, 106(8): 487-491.
- Rolfhus, K.R. and W.F. Fitzgerald. 1995. "Linkages Between Atmospheric Mercury Deposition and the Methylmercury Content of Marine Fish." *Wat. Air Soil Pollut.*, 80: 291-297.
- Rosental, R., G.A. Eagle, and M.J. Orren. 1986. "Trace Metal Distribution in Different Chemical Fractions of Nearshore Marine Sediments." *Estuar. Cstl. Shelf Sci.*, 22: 303-324.
- Ruby, M.V., A. Davis, T.E. Link, R. Schoof, R.L. Chaney, G.B. Freeman, and P. Bergstrom. 1993. "Development of an *In Vitro* Screening Test to Evaluate the *In Vivo* Bioaccessibility of Ingested Mine-Waste Lead." *Environ. Sci. Technol.*, 27(13): 2870-2877.
- Ruby, M.V., A. Davis, R. Schoof, S. Eberle, and C.M. Sellstone. 1996. "Estimation of Lead and Arsenic Bioavailability Using a Physiologically Based Extraction Test." *Environ. Sci. Technol.*, 30(2): 422-430.
- Ruby, M.V., R. Schoof, W. Brattin, M. Goldade, G. Post, M. Harnois, D.E. Mosby, S.W. Casteel, W. Berti, M. Carpenter, D. Edwards, D. Cragin, and W. Chappell. 1999. "Advances in Evaluating the Oral Bioavailability of Inorganics in Soil for Use in Human Health Risk Assessment." *Environ. Sci. Technol.*, 33(21): 3697-3705.
- SAIC, see Science Applications International Corporation.
- Salomons, W., N.M. de Rooij, H. Derdijk, and J. Bril. 1987. "Sediments as a Source for Contaminants?" *Hydrobiol.*, 149: 13-30.
- Salomons, W. and U. Förstner. 1984. *Metals in the Hydrosphere*. Springer-Verlag, Berlin.

- Samant, H.S., K.G. Doe, and O.C. Vaidya. 1990. "An Integrated Chemical and Biological Study of the Bioavailability of Metals in Sediments from Two Contaminated Harbours in New Brunswick, Canada." *Sci. Tot. Environ.*, 96: 253-268.
- Sample, B.E. and G.W. Suter II. 1994. *Estimating Exposure of Terrestrial Wildlife to Contaminants*. ES/ER/TM-125, Oak Ridge National Laboratory.
- Sample, B.E., J.J. Beuchamp, R.A. Efroymsen, G.W. Suter, II, and T.L. Ashwood. 1998. *Development and Validation of Bioaccumulation Models for Earthworms*. Environmental Restoration Program, U.S. Dept. of Energy, Oak Ridge National Laboratory, Oak Ridge, TN. ES/ER/TM-220.
- Schilderman, P.A.E.L., E.J.C. Moonen, P. Kembers, and J.C.S. Kleinjans. 1997. "Bioavailability of Soil-Adsorbed Cadmium in Orally Exposed Male Rats." *Environ. Health Perspect.*, 105: 234-238.
- Schoof, R.A., M.K. Butcher, C. Sellstone, R.W. Ball, J.R. Fricke, V. Keller, and B. Keehn. 1995. "An Assessment of Lead Absorption from Soil Affected by Smelter Emissions." *Environ. Geochem. Health*, 17(4): 189-199.
- Schoof, R.A. and G.B. Freeman. 1995. "Oral Bioavailability of Lead and Cadmium in Soil from a Smelter Site." Poster presented at the Seventh International Congress of Toxicology, Seattle, WA. July 3-6, 1995.
- Schoof, R.A. and J.B. Nielsen. 1997. "Evaluation of Methods for Assessing the Oral Bioavailability of Inorganic Mercury in Soil." *Risk Anal.*, 17(5): 545-555.
- Schafer, H.A., G.P. Hershelman, D.R. Young, and A.J. Mearns. 1982. "Contaminants in Ocean Food Webs." In: W. Bascom (Ed.), *Coastal Water Research Project. Biennial Report for the Years 1981-1982*, pp. 17-28. Southern California Coastal Water Research Project, Long Beach, CA.
- Schropp, S.J., F.G. Lewis, H.L. Windom, J.D. Ryan, F.D. Calder, and L.C. Burney. 1990. "Interpretation of Metal Concentrations in Estuarine Sediments of Florida Using Aluminum as a Reference Element." *Estuaries*, 13: 227-235.
- Science Applications International Corporation. 1994. *East Fork Poplar Creek-Sewer Line Beltway Remedial Investigation Report: Addendum*. DOE/OR/02-1119&D2/A1. Oak Ridge, TN.
- Shea, D. and G.R. Helz. 1988. "The Solubility of Copper in Sulfidic Waters: Sulfide and Polysulfide Complexes in Equilibrium with Covellite." *Geochim. Cosmochim. Acta*, 52: 1815-1825.
- Shea, D. and W.A. MacCrehan. 1988. "Role of Biogenic Thiols in the Solubility of Sulfide Minerals." *Sci. Tot. Environ.*, 73: 135-141.
- Sheppard, S.C., W.G. Evenden, and W.J. Schwartz. 1995. "Heavy Metals in the Environment, Ingested Soil: Bioavailability of Sorbed Lead, Cadmium, Cesium, Iodine, and Mercury." *J. Environ. Qual.*, 24: 498-505.
- Simkiss, K. and M.G. Taylor. 1989. "Metal Fluxes Across the Membranes of Aquatic Organisms." *Rev. Aquat. Sci.*, 1: 173-188.

- Skei, J.M., N.B. Price, S.E. Calvert, and E. Hogdahl. 1972. "The Distribution of Heavy Metals in Sediments of Sörfjord, West Norway." *Water Air Soil Pollut.*, 1: 452-461.
- Smith, D.R., S. Niemeyer, J.A. Estes, and A.R. Flegal. 1990. "Stable Lead Isotopes Evidence Anthropogenic Contamination in Alaska Sea Otters." *Environ. Sci. Technol.*, 24: 1517-1521.
- Snoeij, N.J., A.H. Penninks, and W. Seinen. 1987. "Biological Activity of Organotin Compounds: an Overview." *Environ. Res.*, 44: 335-353.
- Soma, M., A. Tanaka, H. Seyama, and K. Satake. 1994. "Characterization of Arsenic in Lake Sediments by X-Ray Photoelectron Spectroscopy." *Geochim. Cosmochim. Acta*, 58: 2743-2745.
- Stang, P.M. and P.F. Seligman. 1986. "Distribution and Fate of Butyltin Compounds in the Sediment of San Diego Bay." In: *Proceedings of the Oceans 86 Conference: Science-Engineering-Adventure*, pp. 1256-1261. Marine Technology Society, Washington, DC.
- Sullivan, M.F., J.T. Hardy, B.M. Miller, R.L. Buschbom, and T.C. Siewidki. 1984. "Absorption and Distribution of Cadmium in Mice Fed Diets Containing Either Inorganic or Oyster-Incorporated Cadmium." *Toxicol. Appl. Pharmacol.*, 72: 210-217.
- Talmage, S.S. and B.T. Walton. 1993. "Food Chain Transfer and Potential Renal Toxicity to Small Mammals at a Contaminated Terrestrial Field Site." *Ecotoxicology*, 2: 243-256.
- Tessier, A. and P.G.C. Campbell. 1987. "Partitioning of Trace Metals in Sediments: Relationships with Bioavailability." *Hydrobiologia*, 149: 43-52.
- Thompson, D.R. 1990. "Metal Levels in Marine Vertebrates." In: R.W. Furness and P.S. Rainbow (Eds.), *Heavy Metals in the Marine Environment*, pp. 143-182. CRC Press, Boca Raton, FL.
- U.S. Environmental Protection Agency. 1989. *Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation Manual (Part A, Baseline Risk Assessment)*. EPA/540/1-89/002. Office of Emergency and Remedial Response, Washington, DC.
- U.S. Environmental Protection Agency. 1991a. *Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation Manual (Part B, Development of Risk-Based Preliminary Remediation Goals)*. EPA/540/R-92/003. Office of Emergency and Remedial Response, Washington, DC.
- U.S. Environmental Protection Agency. 1991b. *Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation Manual (Part C, Risk Evaluation of Remedial Alternatives)*. EPA/540/R-92/004. Office of Emergency and Remedial Response, Washington, DC.
- U.S. Environmental Protection Agency. 1992. *Dermal Exposure Assessment: Principles and Applications*. EPA/600/8-91/011B. Office of Research and Development, Washington, DC.
- U.S. Environmental Protection Agency. 1994a. *Guidance Manual for the Integrated Exposure Uptake Biokinetic Model for Lead in Children*. EPA/540/R-93/081. Office of Emergency and Remedial Response, Washington, DC.
- U.S. Environmental Protection Agency. 1994b. *Guidance for the Data Quality Objectives Process*. EPA/600/R-96/055. Office of Research and Development, Washington, DC.

- U.S. Environmental Protection Agency. 1996a. *Soil Screening Guidance: Technical Background Document*. EPA/540/R-95/128. Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. Environmental Protection Agency. 1996b. *Bioavailability of Lead in Slag and Soil Samples from the Murray Smelter Superfund Site*. Document Control No. 04800-030-0163. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1996c. *Bioavailability of Lead in Soil Samples from the New Jersey Zinc NPL Site, Palmerton, Pennsylvania*. Document Control No. 04800-030-0159. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1996d. *Bioavailability of Lead in Soil Samples from the Jasper County, Missouri, Superfund Site*. Document Control No. 04800-030-0161. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1996e. *Bioavailability of Lead in Soil Samples from the Smuggler Mountain NPL Site, Aspen, Colorado*. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1998a. *Bioavailability of Lead in Soil and Mine Waste from the California Gulch NPL Site, Leadville, Colorado*. Document Control No. 04800-030-0178. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1998b. *Bioavailability of Lead in a Slag Sample from the Midvale Slag NPL Site, Midvale, Utah*. Document Control No. 04800-030-0166. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1998c. *Bioavailability of Lead in a Soil Sample from the Butte NPL Site, Butte, Montana*. Document Control No. 04800-030-0165. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1998d. *Bioavailability of Lead in Unweathered Galena-Enriched Soil*. Document Control No. 04800-030-0171. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1998e. *Bioavailability of Lead in Paint*. Document Control No. 04800-030-0170. U.S. EPA Region VIII, Denver, CO.
- U.S. Environmental Protection Agency. 1998f. *Guidelines for Ecological Risk Assessment*. EPA/630/R-95/002F. U.S. EPA Risk Assessment Forum, Washington, DC. April.
- U.S. Environmental Protection Agency. 1998g. *Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation Manual, Supplemental Guidance; Dermal Risk Assessment, Interim Guidance*. November 5.
- U.S. Environmental Protection Agency. 1999. *U.S. EPA Region IX Preliminary Remediation Goal Tables*. Available at: www.epa.gov/region09/waste/sfund/prg/. December 3.
- U.S. Environmental Protection Agency. 2000. *U.S. EPA Region III Risk-Based Concentration Table: April 2000 Update*. Available at: www.epa.gov/reg3hwmd/risk/riskmenu.htm. April.
- U.S. EPA, see U.S. Environmental Protection Agency.

- Wahlberg, J.E. and E. Skog. 1963. "The Percutaneous Absorption of Sodium Chromate (^{51}Cr) in the Guinea Pig." *Acta. Derm. Venereol.*, 43: 102-108.
- Wainman, T., R.E. Hazen, and P.J. Liroy. 1994. "The Extractability of Cr(VI) from Contaminated Soil in Synthetic Sweat." *J. Expo. Anal. Environ. Epidemiol.*, 4(2): 171-81.
- Warren, L.J. 1981. "Contamination of Sediments by Lead, Zinc, and Cadmium: a Review." *Environ. Pollut. (Series B)*, 2:401-436.
- Weber, J.H., R. Evans, S.J. Jones, and M.E. Hines. 1998. "Conversion of Mercury (II) into Mercury (0), Monomethylmercury Cation, and Dimethylmercury in Saltmarsh Sediment Slurries." *Chemosphere*, 35: 1669-1687.
- Wen-Xiong, W. and Fisher. 1996. "Assimilation of Trace Elements by the Mussel *Mytilus edulis*: Effects of Diatom Chemical Composition." *Mar. Biol.*, 125: 715-724.
- Wester, R.C., H.I. Maibach, L. Sedik, J. Melendres, S. Dizio, and M. Wade. 1992. "In Vitro Percutaneous Absorption of Cadmium from Water and Soil into Human Skin." *Fund. Appl. Toxicol.*, 19: 1-5.
- Wester, R.C., H.I. Maibach, L. Sedik, J. Melendres, and M. Wade. 1993a. "In Vivo and In Vitro Percutaneous Absorption and Skin Decontamination of Arsenic from Water and Soil." *Fund. Appl. Toxicol.*, 20: 336-340.
- Wester, R.C., D.A.W. Bucks, and H.I. Maibach. 1993b. "Percutaneous Absorption of Contaminants from Soil." In: R.G.M. Wang, J.B. Knaack, and H.I. Maibach (Eds), *Health Risk Assessment: Dermal and Inhalation Exposure and Absorption of Toxicants*, pp. 145-158. CRC Press, Boca Raton, FL.
- Wester, R.C., F. Logan, H. Maibach, M. Wade, and K. Hoang. 1995. "In Vitro Percutaneous Absorption of Mercury from Water and Soil Through Human Skin." *The Toxicologist*, 15(1): 135.
- Witmer, C.M., H.S. Park, and S.I. Shupack. 1989. "Mutagenicity and Disposition of Chromium." *Sci. Total Environ.*, 86: 131-138.
- Witmer, C.M., R. Harris, and S.I. Shupack. 1991. "Oral Bioavailability of Chromium from a Specific Site." *Environ. Health Perspect.*, 92: 105-110.
- Wood, J.M. 1987. "Biological Processes Involved in the Cycling of Elements Between Soil or Sediments and the Aqueous Environment." *Hydrobiol.*, 149: 31-42.
- Wood, T.M., A.M. Baptista, J.S. Kuwabara, and A.R. Flegal. 1995. "Diagnostic Modeling of Trace Metal Partitioning in South San Francisco Bay." *Limnol. Oceanogr.*, 40: 345-358.